

BSDEs under partial information and financial applications.

Claudia Ceci, Alessandra Cretarola, Francesco Russo

▶ **To cite this version:**

Claudia Ceci, Alessandra Cretarola, Francesco Russo. BSDEs under partial information and financial applications.. Stochastic Processes and their Applications, 2014, 10.1016/j.spa.2014.03.003 . hal-00822988

HAL Id: hal-00822988
https://inria.hal.science/hal-00822988

Submitted on 15 May 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BSDES UNDER PARTIAL INFORMATION AND FINANCIAL APPLICATIONS

CLAUDIA CECI, ALESSANDRA CRETAROLA, AND FRANCESCO RUSSO

Abstract. In this paper we provide existence and uniqueness results for the solution of BSDEs driven by a general square integrable martingale under partial information. We discuss some special cases where the solution to a BSDE under restricted information can be derived by that related to a problem of a BSDE under full information. In particular, we provide a suitable version of the Föllmer-Schweizer decomposition of a square integrable random variable working under partial information and we use this achievement to investigate the local risk-minimization approach for a semimartingale financial market model.

1. Introduction

The goal of this paper is to provide existence and uniqueness results for backward stochastic differential equations (in short BSDEs) driven by a general càdlàg square integrable martingale under partial information and to apply such results to provide a financial application.

Frameworks affected by incomplete information represent an interesting issue arising in many problems. Mathematically, this means to consider an additional filtration $\mathbb{H} = (\mathcal{H}_t)_{0\leq t\leq T}$ smaller than the full information flow $\mathbb{F} = (\mathcal{F}_t)_{0\leq t\leq T}$, with T denoting a finite time horizon. A typical example arises when $\mathcal{H}_t = \mathcal{F}_{(t-\tau)^+}$ where $\tau \in (0,T)$ is a fixed delay and $(t-\tau)^+ := \max\{0, t-\tau\}$ with $t \in [0,T]$, or in a financial market where the stock prices can only be observed at discrete time instants or their dynamics depends on an unobservable stochastic factor and $\mathbb{H}$ denotes the information available to investors (see for instance [6], [7], [8], [14], [15]).

For BSDEs driven by a general càdlàg martingale beyond the Brownian setting, there exist very few results in the literature (besides the pioneering work of [4], see [21], [10] and more recently [9], [3] and [5], as far as we are aware). In [9] the authors study for the first time such a general case when there are restrictions on the available information by focusing on BSDEs whose driver is equal to zero. Let $T \in (0,\infty)$ be a fixed time horizon and ξ a square-integrable $\mathcal{F}_T$-measurable random variable which denotes the terminal condition. In this paper we consider general BSDEs of the form:

$$Y_t = \xi + \int_t^T f(s, Y_{s-}, Z_s)\mathrm{d}\langle M\rangle_s - \int_t^T Z_s \mathrm{d}M_s - (O_T - O_t), \quad 0 \leq t \leq T, \tag{1.1}$$

2010 *Mathematics Subject Classification.* 60H10, 60H30, 91B28.

Key words and phrases. Backward stochastic differential equations, partial information, Föllmer-Schweizer decomposition, risk-minimization.

driven by a square-integrable càdlàg $\mathbb{F}$-martingale $M = (M_t)_{0\leq t\leq T}$, with $\mathbb{F}$-predictable quadratic variation $\langle M\rangle = (\langle M\rangle_t)_{0\leq t\leq T}$, where $O = (O_t)_{0\leq t\leq T}$ is a square-integrable $\mathbb{F}$-martingale, satisfying a suitable orthogonality condition that we will make more precise in the next section. The driver of the equation is denoted by f and for each $(y, z) \in \mathbb{R}\times\mathbb{R}$, the process $f(\cdot, \cdot, y, z) = (f(\cdot, t, y, z))_{0\leq t\leq T}$ is $\mathbb{F}$-predictable.

We look for a solution (Y, Z) to equation (1.1) under partial information, where $Y = (Y_t)_{0\leq t\leq T}$ is a càdlàg $\mathbb{F}$-adapted process such that $\mathbb{E}\left[\sup_{0\leq t\leq T}|Y_t|^2\right] < \infty$ and $Z = (Z_t)_{0\leq t\leq T}$ is an $\mathbb{H}$-predictable process such that $\mathbb{E}\left[\int_0^T |Z_s|^2 \mathrm{d}\langle M\rangle_s\right] < \infty$.

Our first important achievement, stated in Theorem 2.12, concerns existence and uniqueness properties of the solution to such BSDEs. We get such results by assuming f uniformly Lipschitz with respect to (y, z) and the behavior of $\langle M\rangle$ to be controlled by a deterministic function. Moreover, we provide in Proposition 2.14 a representation of the solution to BSDEs under restricted information in terms of the Radon-Nikodým derivative of two $\mathbb{H}$-predictable dual projections involving the solution of a problem under full information. Thanks to this result, in the particular case where the driver f does not depend on z, we give in Proposition 2.16 an explicit characterization of the solution to BSDEs under restricted information in terms of the solution to the corresponding BSDEs under full information. Finally, as an illustrative example, we discuss the special case of delayed information, that is, when $\mathcal{H}_t = \mathcal{F}_{(t-\tau)^+}$ for each $t \in [0, T]$, with $\tau \in (0, T)$ being fixed, once we assume that $\langle M\rangle$ and $f(\cdot, \cdot, y, z)$ are $\mathbb{H}$-predictable processes and f does not depend on y. Proposition 2.18 ensures existence of the solution to the BSDEs under restricted information by a constructive procedure under weaker conditions on f with respect to the general theorem.

As a financial application we discuss the local risk-minimization approach for partially observable semimartingale models. The local risk-minimization approach is a quadratic hedging method for contingent claims in incomplete markets which keeps the replication constraint and looks for a hedging strategy (in general not self-financing) with minimal cost, see e.g. [12] and [23] for a further discussion on this issue. The study of this approach under partial information in full generality is still an interesting topic to discuss. The first step was done by [12], where they complete the information starting from the reference filtration and recover the optimal strategy by means of predictable projections with respect to the enlarged filtration. Some further contributions in this direction can be found in [22] and [9] in the case where the underlying price process is a (local) martingale under the real-world probability measure. In [22], the author provides an explicit expression for risk-minimizing hedging strategies under restricted information in terms of predictable dual projections, whereas in [9], by proving a version of the Galtchouk-Kunita-Watanabe decomposition that works under partial information, the authors extend the results of [13] to the partial information framework and show how their result fits in the approach of [22]. Furthermore, an application of the local risk-minimization approach in the case of incomplete information to defaultable markets in the sense of [12] can be found in [2].

Here, we consider a more general situation since we allow the underlying price process to be represented by a semimartingale under the real-world probability measure. More precisely, in Proposition 3.10 we provide a version of the Föllmer-Schweizer decomposition of a square-integrable random variable (that typically represents the payoff of a contract) with respect to the underlying price process, that works under partial information.

Then, we study the relationship between the Föllmer-Schweizer decomposition of a contingent claim under partial information and the existence of a locally risk-minimizing strategy according to the partial information framework.

In addition, we discuss the case where the underlying price process can exhibit jumps in the classical full information setting.

The paper is organized as follows. In Section 2 we formulate the problem for BSDEs under partial information, we prove existence and uniqueness properties of solutions and we give the representation results in terms of $\mathbb{H}$-predictable dual projections. Section 2 concludes with a discussion of some special cases. Section 3 is devoted to the study of local risk-minimization under partial information via BSDEs. A discussion about the case of complete information in presence of jumps in the underlying price process can be found in Section 3.1. Finally, some detailed definitions and technical results are gathered in Section A in Appendix.

2. Backward stochastic differential equations under partial information

Let us fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ endowed with a filtration $\mathbb{F} := (\mathcal{F}_t)_{0\leq t\leq T}$, where $\mathcal{F}_t$ represents the full information at time t and T denotes a fixed and finite time horizon. We assume that $\mathcal{F} = \mathcal{F}_T$. Then we consider a subfiltration $\mathbb{H} := (\mathcal{H}_t)_{0\leq t\leq T}$ of $\mathbb{F}$, i.e. $\mathcal{H}_t \subseteq \mathcal{F}_t$, for each $t \in [0, T]$, corresponding to the available information level. We remark that both filtrations are assumed to satisfy the usual hypotheses of completeness and right-continuity, see e.g. [17].
For simplicity we only consider the one-dimensional case. Extensions to several dimensions are straightforward and left to the reader. The data of the problem are:

- an $\mathbb{R}$-valued square-integrable (càdlàg) $\mathbb{F}$-martingale $M = (M_t)_{0\leq t\leq T}$ with $\mathbb{F}$-predictable quadratic variation process denoted by $\langle M\rangle = (\langle M, M\rangle)_{0\leq t\leq T}$;
- a terminal condition $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$[1];
- a coefficient $f : \Omega \times [0, T] \times \mathbb{R} \times \mathbb{R} \longrightarrow \mathbb{R}$, such that, for each $(y, z) \in \mathbb{R} \times \mathbb{R}$, the process $f(\cdot, \cdot, y, z) = (f(\cdot, t, y, z))_{0\leq t\leq T}$ is $\mathbb{F}$-predictable. The random function f is said to be the *driver* of the equation.

We make the following assumptions on the coefficient f.

[1]The space $L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$ denotes the set of all real-valued $\mathcal{F}_T$-measurable random variables H such that $\mathbb{E}\left[|H|^2\right] = \int_\Omega |H|^2 \mathrm{d}\mathbb{P} < \infty$.

Assumption 2.1.

(i) *f is uniformly Lipschitz with respect to (y,z): there exists a constant $K \geq 0$ such that for every $(y,z),(y',z') \in \mathbb{R}\times\mathbb{R}$,*

$$|f(\omega,t,y,z) - f(\omega,t,y',z')| \leq K\left(|y-y'| + |z-z'|\right), \quad (\mathbb{P}\otimes\langle M\rangle) - a.e.\ on\ \Omega\times[0,T];$$

(ii) *the following integrability condition is satisfied:*

$$\mathbb{E}\left[\int_0^T |f(t,0,0)|^2 \mathrm{d}\langle M\rangle_t\right] < \infty.$$

To describe the parameters and the solution of BSDEs, we introduce the following spaces:

- $\mathcal{S}^2_{\mathcal{F}}(0,T)$, the set of all càdlàg $\mathbb{F}$-adapted processes $\phi = (\phi_t)_{0\leq t\leq T}$ such that $\|\phi\|^2_{\mathcal{S}^2} := \mathbb{E}\left[\sup_{0\leq t\leq T}|\phi_t|^2\right] < \infty$;
- $\mathcal{M}^2_{\mathcal{H}}(0,T)$ ($\mathcal{M}^2_{\mathcal{F}}(0,T)$), the set of all $\mathbb{H}$-predictable (respectively $\mathbb{F}$-predictable) processes $\varphi = (\varphi_t)_{0\leq t\leq T}$ such that $\|\varphi\|^2_{\mathcal{M}^2} := \mathbb{E}\left[\int_0^T |\varphi_s|^2 \mathrm{d}\langle M\rangle_s\right] < \infty$;
- $\mathcal{L}^2_{\mathcal{F}}(0,T)$, the set of all $\mathbb{F}$-martingales $\psi = (\psi_t)_{0\leq t\leq T}$ with $\psi_0 = 0$, such that $\|\psi\|^2_{\mathcal{L}^2} := \mathbb{E}\left[\langle\psi\rangle_T\right] = \mathbb{E}\left[\psi_T^2\right] < \infty$.

We now give the definitions of solution in a full and in a partial information framework, respectively.

Definition 2.2. *A solution of the BSDE*

$$\tilde{Y}_t = \xi + \int_t^T f(s,\tilde{Y}_{s-},\tilde{Z}_s)\mathrm{d}\langle M\rangle_s - \int_t^T \tilde{Z}_s \mathrm{d}M_s - (\tilde{O}_T - \tilde{O}_t), \quad 0\leq t\leq T, \tag{2.1}$$

with data (ξ, f) under complete information, is a triplet $(\tilde{Y},\tilde{Z},\tilde{O}) = (\tilde{Y}_t,\tilde{Z}_t,\tilde{O}_t)_{0\leq t\leq T}$ of processes with values in $\mathbb{R}\times\mathbb{R}\times\mathbb{R}$ satisfying (2.1), such that

$$(\tilde{Y},\tilde{Z},\tilde{O}) \in \mathcal{S}^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{F}}(0,T)\times\mathcal{L}^2_{\mathcal{F}}(0,T),$$

where $\tilde{O}$ is strongly orthogonal to M (i.e. $\langle\tilde{O},M\rangle_t = 0$ $\mathbb{P}$-a.s., for every $t\in[0,T]$).

Definition 2.3. *A solution of the BSDE*

$$Y_t = \xi + \int_t^T f(s,Y_{s-},Z_s)\mathrm{d}\langle M\rangle_s - \int_t^T Z_s \mathrm{d}M_s - (O_T - O_t), \quad 0\leq t\leq T, \tag{2.2}$$

with data $(\xi, f, \mathbb{H})$ under partial information, is a triplet $(Y,Z,O) = (Y_t,Z_t,O_t)_{0\leq t\leq T}$ of processes with values in $\mathbb{R}\times\mathbb{R}\times\mathbb{R}$ satisfying (2.2), such that

$$(Y,Z,O) \in \mathcal{S}^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{H}}(0,T)\times\mathcal{L}^2_{\mathcal{F}}(0,T),$$

where O satisfies the orthogonality condition

$$\mathbb{E}\left[O_T\int_0^T \varphi_t \mathrm{d}M_t\right] = 0, \tag{2.3}$$

for all processes $\varphi\in\mathcal{M}^2_{\mathcal{H}}(0,T)$.

Remark 2.4. *Sometimes in the literature, only the couple (Y,Z) identifies the solution of a BSDE of the form (2.1) or (2.2). Indeed, this is reasonable since the $\mathbb{F}$-martingale O is uniquely determined by the processes Y and Z that satisfy the equation.*

Remark 2.5. *The orthogonality condition given in* (2.3) *is weaker than the classical strong orthogonality condition considered in Definition 2.2. Indeed, set* $N_t = \int_0^t \varphi_s \mathrm{d}M_s$, *for each* $t \in [0,T]$, *where* $\varphi \in \mathcal{M}^2_{\mathcal{H}}(0,T)$. *If* $\psi \in \mathcal{L}^2_{\mathcal{F}}(0,T)$ *is such that*

$$\langle \psi, M\rangle_t = 0 \quad \mathbb{P}-a.s., \quad \forall t \in [0,T],$$

then

$$\langle \psi, N\rangle_t = \int_0^t \varphi_s \mathrm{d}\langle \psi, M\rangle_s = 0 \quad \mathbb{P}-a.s., \quad \forall t \in [0,T].$$

Consequently, ψN *is an* $\mathbb{F}$*-martingale null at zero, that implies*

$$\mathbb{E}\left[\psi_t N_t\right] = 0, \quad \forall t \in [0,T],$$

and in particular condition (2.3).

Remark 2.6. *Let* $\psi \in \mathcal{L}^2_{\mathcal{F}}(0,T)$. *Since for any* $\mathbb{H}$*-predictable process* φ, *the process* $\mathbf{1}_{(0,t]}(s)\varphi_s$, *with* $t \leq T$, *is* $\mathbb{H}$*-predictable, condition* (2.3) *implies that for every* $t \in [0,T]$ *and for each* $\varphi \in \mathcal{M}^2_{\mathcal{H}}(0,T)$, *we have*

$$\mathbb{E}\left[\psi_T \int_0^t \varphi_s \mathrm{d}M_s\right] = 0.$$

Then, by conditioning with respect to $\mathcal{F}_t$ *(note that* ψ *is an* $\mathbb{F}$*-martingale), for every* $\varphi \in \mathcal{M}^2_{\mathcal{H}}(0,T)$, *we get*

$$\mathbb{E}\left[\psi_t \int_0^t \varphi_s \mathrm{d}M_s\right] = \mathbb{E}\left[\int_0^t \varphi_s \mathrm{d}\langle M, \psi\rangle_s\right] = 0 \quad \forall t \in [0,T].$$

From this last equality, we can argue that in the case of full information, i.e., when $\mathcal{H}_t = \mathcal{F}_t$, *for each* $t \in [0,T]$, *condition* (2.3) *is equivalent to the strong orthogonality condition between* ψ *and* M *(see e.g. Lemma 2 and Theorem 36, Chapter IV, page 180 of [17] for a rigorous proof).*

In the sequel, we will say that a square-integrable $\mathbb{F}$-martingale O is *weakly orthogonal* to M if condition (2.3) holds for all processes $\varphi \in \mathcal{M}^2_{\mathcal{H}}(0,T)$.

2.1. **Existence and Uniqueness.** Our aim is to investigate existence and uniqueness of solutions to the BSDE (2.2) with data $(\xi, f, \mathbb{H})$ driven by the general martingale M in the sense of Definition 2.3. The case $f \equiv 0$ in (2.2), has been studied in [9], where a key role is played by the Galtchouk-Kunita-Watanabe decomposition under partial information that we recall here for reader's convenience.

Proposition 2.7. *Let* $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$. *There exists a unique decomposition of the form*

$$\xi = U_0 + \int_0^T H_t^{\mathcal{H}} \mathrm{d}M_t + A_T, \quad \mathbb{P}-a.s., \tag{2.4}$$

where $U_0 \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; \mathbb{R})$, $H^{\mathcal{H}} = (H_t^{\mathcal{H}})_{0\leq t\leq T} \in \mathcal{M}^2_{\mathcal{H}}(0,T)$ *and* $A = (A_t)_{0\leq t\leq T} \in \mathcal{L}^2_{\mathcal{F}}(0,T)$ *weakly orthogonal to* M.

Inspired by [3], we make the following assumption on the $\mathbb{F}$-predictable quadratic variation $\langle M\rangle$ of M.

Assumption 2.8. *There exists a deterministic function $\rho : \mathbb{R}_+ \to \mathbb{R}_+$ with $\rho(0^+) = 0$ such that, $\mathbb{P}$-a.s.,*

$$\langle M\rangle_t - \langle M\rangle_s \leq \rho(t-s), \quad \forall 0 \leq s \leq t \leq T.$$

Example 2.9. *On the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ let us consider a standard Brownian motion W and an independent Poisson random measure $N(\mathrm{d}\zeta, \mathrm{d}t)$ on $Z \times [0,T]$ with non-negative intensity $\nu(\mathrm{d}\zeta)\mathrm{d}t$, where $\nu(\mathrm{d}\zeta)$ is a σ-finite measure on a measurable space $(Z, \mathcal{Z})$. Denote by $\tilde{N}$ the corresponding compensated measure defined by*

$$\tilde{N}(\mathrm{d}\zeta, \mathrm{d}t) = N(\mathrm{d}\zeta, \mathrm{d}t) - \nu(\mathrm{d}\zeta)\mathrm{d}t.$$

Let M be given by

$$M_t = M_0 + \int_0^t \bar{\sigma}_s \mathrm{d}W_s + \int_0^t \int_Z \bar{K}(\zeta; s)\tilde{N}(\mathrm{d}\zeta, \mathrm{d}s), \quad t \in [0,T],$$

with $\bar{\sigma} = (\bar{\sigma}_t)_{0\leq t\leq T}$ and $\bar{K} = (\bar{K}(\cdot; t))_{0\leq t\leq T}$ being $\mathbb{R}$-valued, $\mathbb{F}$-adapted and $\mathbb{F}$-predictable processes respectively, and satisfying

$$\mathbb{E}\left[\int_0^T \bar{\sigma}_s^2 \mathrm{d}s + \int_0^T \int_Z \bar{K}^2(\zeta; s)\nu(\mathrm{d}\zeta)\mathrm{d}s\right] < \infty.$$

Then, M is a square-integrable $\mathbb{F}$-martingale with $\mathbb{F}$-predictable quadratic variation process $\langle M\rangle$ given by

$$\langle M\rangle_t = \int_0^t \left(\bar{\sigma}_s^2 + \int_Z \bar{K}^2(\zeta; s)\nu(\mathrm{d}\zeta)\right)\mathrm{d}s, \quad t \in [0,T].$$

If in addition we assume that there exists a positive constant $\bar{C}$ such that

$$\bar{\sigma}_t^2 + \int_Z \bar{K}^2(\zeta; t)\nu(\mathrm{d}\zeta) \leq \bar{C} \quad \mathrm{d}\mathbb{P} \times \mathrm{d}t - a.e. \tag{2.5}$$

then Assumption 2.8 is fulfilled with $\rho(t-s) = \bar{C}(t-s)$, with $0 \leq s \leq t \leq T$.

Let us observe that in particular condition (2.5) is satisfied if both processes $\bar{\sigma}$ and $\bar{K}$ are bounded and $\nu(\{\zeta \in Z : \bar{K}(\zeta; t) \neq 0\}) < \infty$ for every $t \in [0,T]$.

We start with the following lemma.

Lemma 2.10. *Let Assumption 2.1 hold and assume that $\langle M\rangle_T \leq C(T)$ $\mathbb{P}$-a.s., where $C(T)$ is a positive constant depending on T. Let $(U,V) = (U_t, V_t)_{0\leq t\leq T} \in \mathcal{S}^2_{\mathcal{F}}(0,T) \times \mathcal{M}^2_{\mathcal{H}}(0,T)$. Then the BSDE*

$$Y_t = \xi + \int_t^T f(s, U_{s-}, V_s)\mathrm{d}\langle M\rangle_s - \int_t^T Z_s \mathrm{d}M_s - (O_T - O_t), \quad 0 \leq t \leq T, \tag{2.6}$$

has a solution with data $(\xi, f, \mathbb{H})$ under partial information in the sense of Definition 2.3.

Proof. Firstly, we set

$$Y_t = \mathbb{E}\left[\xi + \int_t^T f(s, U_{s-}, V_s)\mathrm{d}\langle M\rangle_s \middle| \mathcal{F}_t\right], \quad t \in [0,T].$$

Here Y is a càdlàg $\mathbb{F}$-adapted process and moreover

$$|Y_t| \leq m_t := \mathbb{E}\left[|\xi| + \int_0^T |f(s, U_{s-}, V_s)| \mathrm{d}\langle M\rangle_s \middle| \mathcal{F}_t\right], \quad t \in [0,T],$$

where $m = (m_t)_{0\leq t\leq T}$ is a square-integrable $\mathbb{F}$-martingale. Thus, Doob's inequality and Jensen's inequality yield

$$\begin{aligned}
\mathbb{E}\left[\sup_{0\leq t\leq T} |Y_t|^2\right] &\leq \mathbb{E}\left[\sup_{0\leq t\leq T} |m_t|^2\right] \\
&\leq 4 \sup_{0\leq t\leq T} \mathbb{E}\left[|m_t|^2\right] = 4\mathbb{E}\left[\left(|\xi| + \int_0^T |f(s, U_{s-}, V_s)| \mathrm{d}\langle M\rangle_s\right)^2\right] \\
&\leq 8\mathbb{E}\left[|\xi|^2\right] + 8\mathbb{E}\left[\left(\int_0^T |f(s, U_{s-}, V_s)| \mathrm{d}\langle M\rangle_s\right)^2\right].
\end{aligned}$$

By Cauchy-Schwarz inequality and boundedness of $\langle M\rangle$, we get

$$\mathbb{E}\left[\sup_{0\leq t\leq T} |Y_t|^2\right] \leq 8\mathbb{E}\left[|\xi|^2\right] + 8C(T)\mathbb{E}\left[\int_0^T |f(s, U_{s-}, V_s)|^2 \mathrm{d}\langle M\rangle_s\right].$$

Finally, by Assumption 2.1, we have

$$\begin{aligned}
\mathbb{E}\left[\sup_{0\leq t\leq T} |Y_t|^2\right] \leq 8\mathbb{E}\left[|\xi|^2\right] &+ 32C(T)K^2\mathbb{E}\left[\int_0^T (|U_{s-}|^2 + |V_s|^2) \mathrm{d}\langle M\rangle_s\right] \\
&+ 16C(T)\mathbb{E}\left[\int_0^T |f(s,0,0)|^2 \mathrm{d}\langle M\rangle_s\right].
\end{aligned}$$

The right-hand side of previous inequality is finite in view of hypotheses on (U, V), Assumptions 2.1 and 2.8. Hence, $Y \in \mathcal{S}^2_{\mathcal{F}}(0,T)$.

By Proposition 2.7, the square-integrable $\mathcal{F}_T$-measurable random variable

$$\xi + \int_0^T f(s, U_{s-}, V_s) \mathrm{d}\langle M\rangle_s$$

admits a unique Galtchouk-Kunita-Watanabe decomposition under partial information. Setting $Z_t = H^{\mathcal{H}}_t$ and $O_t = A_t$ for every $t \in [0,T]$, see (2.4), this ensures uniqueness of the process $Z \in \mathcal{M}^2_{\mathcal{H}}(0,T)$ and of the process $O \in \mathcal{L}^2_{\mathcal{F}}(0,T)$ satisfying the orthogonality condition (2.3), which verify the BSDE (2.6).

Indeed, taking the conditional expectation with respect to $\mathcal{F}_t$ yields the following identity:

$$\begin{aligned}
\mathbb{E}\left[\xi + \int_0^T f(s, U_{s-}, V_s) \mathrm{d}\langle M\rangle_s \middle| \mathcal{F}_t\right] &= \mathbb{E}\left[U_0 + \int_0^T H^{\mathcal{H}}_s \mathrm{d}M_s + A_T \middle| \mathcal{F}_t\right] \\
&= U_0 + \int_0^t H^{\mathcal{H}}_s \mathrm{d}M_s + A_t \\
&= Y_0 + \int_0^t H^{\mathcal{H}}_s \mathrm{d}M_s + A_t, \quad 0 \leq t \leq T. \qquad (2.7)
\end{aligned}$$

By (2.1) and (2.7) we have that

$$Y_t + \int_0^t f(s, U_{s-}, V_s) \mathrm{d}\langle M\rangle_s = Y_0 + \int_0^t H^{\mathcal{H}}_s \mathrm{d}M_s + A_t, \quad 0 \leq t \leq T,$$

from which we deduce that

$$Y_t = \xi + \int_t^T f(s, U_{s-}, V_s)\mathrm{d}\langle M\rangle_s - \int_t^T H_s^{\mathcal{H}}\mathrm{d}M_s - (A_T - A_t), \quad 0 \le t \le T.$$

□

We keep on the study by giving an estimation result.

Proposition 2.11. *Under Assumptions 2.1 and 2.8, let* (Y, Z, O) (*respectively* (Y', Z', O')) *be a solution of the BSDE* (2.2) *with data* $(\xi, f, \mathbb{H})$ (*respectively with data* $(\xi', f, \mathbb{H})$) *associated to* $(U, V) \in \mathcal{S}^2_{\mathcal{F}}(0,T) \times \mathcal{M}^2_{\mathcal{H}}(0,T)$ (*respectively* $(U', V') \in \mathcal{S}^2_{\mathcal{F}}(0,T) \times \mathcal{M}^2_{\mathcal{H}}(0,T)$). *Then, for each* $0 \le u \le v \le T$, *we have*

$$\begin{aligned}\mathbb{E}\left[\sup_{u\le t\le v}|\delta Y_t|^2 + \int_u^v |\delta Z_s|^2\mathrm{d}\langle M\rangle_s + \langle\delta O\rangle_v - \langle\delta O\rangle_u\right] \\ \le 42\mathbb{E}\left[|\delta Y_v|^2\right] + C(v-u)\mathbb{E}\left[\sup_{u\le t\le v}|\delta U_t|^2 + \int_u^v |\delta V_s|^2\mathrm{d}\langle M\rangle_s\right],\end{aligned} \tag{2.8}$$

where $C(r) = 42K^2\max\{\rho^2(r), \rho(r)\}$ *and* δY *stands for* $Y - Y'$ *and so on.*

Proof. For reader's convenience, here we provide briefly the proof of (2.8). It is formally analogous to the one of Proposition 7 in [3]. The difference is due to the orthogonality condition we consider in this framework. We start by the following equation: for every $t \in [0, v] \subseteq [0, T]$, set

$$\delta Y_t = \delta Y_v + \int_t^v \left(f(s, U_{s-}, V_s) - f(s, U'_{s-}, V'_s)\right)\mathrm{d}\langle M\rangle_s - \int_t^v \delta Z_s\mathrm{d}M_s - (\delta O_v - \delta O_t). \tag{2.9}$$

Since f is K-Lipschitz in virtue of Assumption 2.1, for any $t \in [0, v]$ we have

$$|\delta Y_t| \le \mathbb{E}\left[|\delta Y_v| + K\int_t^v (|\delta U_{s-}| + |\delta V_s|)\,\mathrm{d}\langle M\rangle_s \middle| \mathcal{F}_t\right] \le \tilde{m}_t,$$

where $\tilde{m} = (\tilde{m}_t)_{0\le t\le T}$, defined by $\tilde{m}_t := \mathbb{E}\left[|\delta Y_v| + K\int_u^v (|\delta U_{s-}| + |\delta V_s|)\,\mathrm{d}\langle M\rangle_s \middle| \mathcal{F}_t\right]$ for each $t \in [0, T]$, is a square-integrable $\mathbb{F}$-martingale. Doob's inequality gives

$$\begin{aligned}\mathbb{E}\left[\sup_{u\le t\le v}|\delta Y_t|^2\right] &\le \mathbb{E}\left[\sup_{u\le t\le v}|\tilde{m}_t|^2\right] \le 4\sup_{u\le t\le v}\mathbb{E}\left[|\tilde{m}_t|^2\right] \\ &\le 4\mathbb{E}\left[\left(|\delta Y_v| + K\int_u^v (|\delta U_{s-}| + |\delta V_s|)\,\mathrm{d}\langle M\rangle_s\right)^2\right].\end{aligned} \tag{2.10}$$

Furthermore, since δO satisfies the orthogonality condition (2.3), it is easy to check that

$$\mathbb{E}\left[(\delta O_v - \delta O_u)\int_u^v \delta Z_s\mathrm{d}M_s\right] = 0, \quad 0 \le u \le v \le T;$$

then

$$\mathbb{E}\left[\int_u^u |\delta Z_s|^2\mathrm{d}\langle M\rangle_s + \langle\delta O\rangle_v - \langle\delta O\rangle_u\right] = \mathbb{E}\left[\left|\int_u^v \delta Z_s\mathrm{d}M_s + \delta O_v - \delta O_u\right|^2\right]. \tag{2.11}$$

Hence, taking (2.9) into account we derive

$$\int_u^v \delta Z_s \mathrm{d}M_s + \delta O_v - \delta O_u = \delta Y_v - \delta Y_u + \int_u^v \Big(f(s, U_{s-}, V_s) - f(s, U'_{s-}, V'_s) \Big) \mathrm{d}\langle M \rangle_s.$$

Using the fact that f is K-Lipschitz in virtue of Assumption 2.1, we obtain

$$\left| \int_u^v \delta Z_s \mathrm{d}M_s + \delta O_v - \delta O_u \right| \leq |\delta Y_v| + \sup_{u \leq t \leq v} |\delta Y_t| + K \int_u^v \left(|\delta U_{s-}| + |\delta V_s| \right) \mathrm{d}\langle M \rangle_s.$$

Since (2.10) also holds for $\mathbb{E}\left[\left(\sup_{u \leq t \leq v} |\delta Y_t| \right)^2 \right]$, from the estimate (2.10) and relationship (2.11), we get

$$\begin{aligned} &\mathbb{E}\left[\sup_{u \leq t \leq v} |\delta Y_t|^2 + \int_u^v |\delta Z_s|^2 \mathrm{d}\langle M \rangle_s + \langle \delta O \rangle_v - \langle \delta O \rangle_u \right] \\ &\qquad \leq 14 \mathbb{E}\left[\left(|\delta Y_v| + K \int_u^v \left(|\delta U_{s-}| + |\delta V_s| \right) \mathrm{d}\langle M \rangle_s \right)^2 \right]. \end{aligned}$$

Cauchy-Schwarz inequality together with Assumption 2.8 lead to the estimate

$$\begin{aligned} &\mathbb{E}\left[\sup_{u \leq t \leq v} |\delta Y_t|^2 + \int_u^v |\delta Z_s|^2 \mathrm{d}\langle M \rangle_s + \langle \delta O \rangle_v - \langle \delta O \rangle_u \right] \\ &\qquad \leq 42 \mathbb{E}\left[|\delta Y_v|^2 \right] + C(v-u) \mathbb{E}\left[\sup_{u \leq t \leq v} |\delta U_t|^2 + \int_u^v |\delta V_s|^2 \mathrm{d}\langle M \rangle_s \right], \end{aligned}$$

with $C(v-u) = 42K^2 \max\{\rho^2(v-u), \rho(v-u)\}$. □

Note that, since $\lim_{r \to 0^+} \rho(r) = 0$ by Assumption 2.8, there exists $r_0 \in (0, T)$ such that $42K^2 \max\{\rho^2(v-u), \rho(v-u)\} \leq \frac{1}{6}$ as soon as $r \leq r_0$. Similarly to [3], we introduce the following norm on $\mathcal{S}^2_{\mathcal{F}}(0,T) \times \mathcal{M}^2_{\mathcal{H}}(0,T) \times \mathcal{L}^2_{\mathcal{F}}(0,T)$:

$$\|(Y,Z,O)\|_p^2 := \sum_{k=0}^{\hat{m}-1} (5 \cdot 42)^k \mathbb{E}\left[\sup_{I_k} |Y_t|^2 + \int_{I_k} |Z_s|^2 \mathrm{d}\langle M \rangle_s + \langle O \rangle_{\frac{(k+1)T}{\hat{m}}} - \langle O \rangle_{\frac{kT}{\hat{m}}} \right],$$

where $\hat{m} = [T/r_0] + 1$ is fixed and $I_k = [kT/\hat{m}, (k+1)T/\hat{m}]$, for $0 \leq k \leq \hat{m} - 1$, are $\hat{m}$ intervals that constitute a regular partition of $[0, T]$. This norm is equivalent to the classical one since we have

$$\|(Y,Z,O)\|^2 := \|Y\|^2_{\mathcal{S}^2} + \|Z\|^2_{\mathcal{M}^2} + \|O\|^2_{\mathcal{L}^2} \leq \|(Y,Z,O)\|_p^2 \leq \hat{m}(5 \cdot 42)^{\hat{m}-1} \|(Y,Z,O)\|^2.$$

Thanks to the estimate of Proposition 2.11 and a straightforward computation, we can show that if (Y,Z,O) and (Y',Z',O') are the solutions to the BSDE (2.6) with (ξ, U, V) and (ξ', U', V') respectively, then we have

$$\|(\delta Y, \delta Z, \delta O)\|_p^2 \leq \frac{1}{5} \|(\delta Y, \delta Z, \delta O)\|_p^2 + C(T/\hat{m}) \|(\delta U, \delta V, 0)\|_p^2.$$

Hence

$$\|(\delta Y, \delta Z, \delta O)\|_p^2 \leq \frac{1}{4} \|(\delta U, \delta V, 0)\|_p^2. \tag{2.12}$$

Theorem 2.12. *Let Assumptions 2.1 and 2.8 hold. Given data $(\xi, f, \mathbb{H})$, there exists a unique triplet (Y,Z,O) which solves the BSDE (2.2) under partial information in the sense of Definition 2.3.*

Proof. The idea is to use a fixed point argument. Let us consider the application Φ from $\mathcal{S}^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{H}}(0,T)\times\mathcal{L}^2_{\mathcal{F}}(0,T)$ into itself which is defined by setting $\Phi(U,V,L)=(Y,Z,O)$ where (Y,Z,O) is the solution to the BSDE (2.6). Note that L does not appear and the application is well-defined thanks to Lemma 2.10 and since estimate (2.12) ensures the existence of a unique solution, in the space $\mathcal{S}^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{H}}(0,T)\times\mathcal{L}^2_{\mathcal{F}}(0,T)$, to the BSDE (2.6), once the pair $(U,V)\in\mathcal{S}^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{H}}(0,T)$ is fixed.
Indeed, the estimate (2.12) says that Φ is a contraction with constant $\frac{1}{2}$ if we use the equivalent norm $\|\cdot\|_p$ instead of the the classical one $\|\cdot\|$ on the Banach space $\mathcal{S}^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{H}}(0,T)\times\mathcal{L}^2_{\mathcal{F}}(0,T)$. □

Since it will be useful in the sequel, we recall for reader's convenience the definition of $\mathbb{H}$-predictable dual projection.

Definition 2.13. *Let $G=(G_t)_{0\le t\le T}$ be a càdlàg $\mathbb{F}$-adapted process of integrable variation. The $\mathbb{H}$-predictable dual projection of G is the unique $\mathbb{H}$-predictable process $G^{\mathbb{H}}=(G^{\mathbb{H}}_t)_{0\le t\le T}$ of integrable variation such that*

$$\mathbb{E}\left[\int_0^T\varphi_s\mathrm{d}G^{\mathbb{H}}_t\right]=\mathbb{E}\left[\int_0^T\varphi_s\mathrm{d}G_t\right],$$

for every $\mathbb{H}$-predictable (bounded) process φ.

It is possible to show that BSDEs under partial information can be reduced to full information problems, which however are not described by a BSDE, unless the driver does not depend on z (see Proposition 2.16 below). More precisely, we have the following result.

Proposition 2.14. *Let $(\tilde{Y},\tilde{Z},\tilde{O})\in\mathcal{S}^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{F}}(0,T)\times\mathcal{L}^2_{\mathcal{F}}(0,T)$ be a solution to the problem under complete information*

$$\tilde{Y}_t=\xi+\int_t^T f(s,\tilde{Y}_{s-},\hat{Z}_s)\mathrm{d}\langle M\rangle_s-\int_t^T\tilde{Z}_s\mathrm{d}M_s-(\tilde{O}_T-\tilde{O}_t),\quad 0\le t\le T,\tag{2.13}$$

where $\tilde{O}$ is strongly orthogonal to M and

$$\hat{Z}_t=\frac{\mathrm{d}L^{\mathbb{H}}_t}{\mathrm{d}\langle M\rangle^{\mathbb{H}}_t}\quad L_t:=\int_0^t\tilde{Z}_s\mathrm{d}\langle M\rangle_s,\quad 0\le t\le T.$$

Then the triplet

$$(Y,Z,O)=\left(\tilde{Y},\hat{Z},\tilde{O}+B\right),$$

where $B=\int(\tilde{Z}_s-\hat{Z}_s)\mathrm{d}M_s$ is a square-integrable $\mathbb{F}$-martingale weakly orthogonal to M, is a solution to the BSDE (2.2) under partial information.

Proof. First let us observe that by Proposition 4.8 of [9], $L^{\mathbb{H}}:=(\int\tilde{Z}_s\mathrm{d}\langle M\rangle_s)^{\mathbb{H}}$ is absolutely continuous with respect to $\langle M\rangle^{\mathbb{H}}$, hence $\hat{Z}$ is well defined. By (2.13) we get

$$\tilde{Y}_t=\xi+\int_t^T f(s,\tilde{Y}_{s-},\hat{Z}_s)\mathrm{d}\langle M\rangle_s-\int_t^T\hat{Z}_s\mathrm{d}M_s-\int_t^T(\tilde{Z}_s-\hat{Z}_s)\mathrm{d}M_s-(\tilde{O}_T-\tilde{O}_t),\quad 0\le t\le T.$$

Set $B_t = \int_0^t (\tilde{Z}_s - \hat{Z}_s)\mathrm{d}M_s$, for each $t \in [0,T]$. It is sufficient to prove that $B = (B_t)_{0\leq t\leq T}$ is a square-integrable $\mathbb{F}$-martingale weakly orthogonal to M, that is, for every $\varphi \in \mathcal{M}^2_{\mathcal{H}}(0,T)$ we have

$$\mathbb{E}\left[\int_0^T \varphi_s \mathrm{d}M_s \int_0^T (\tilde{Z}_s - \hat{Z}_s)\mathrm{d}M_s\right] = \mathbb{E}\left[\int_0^T \varphi_s(\tilde{Z}_s - \hat{Z}_s)\langle M\rangle_s\right] = 0.$$

In fact

$$\mathbb{E}\left[\int_0^T \varphi_s \tilde{Z}_s \langle M\rangle_s\right] = \mathbb{E}\left[\int_0^T \varphi_s (\tilde{Z}\langle M\rangle)^{\mathbb{H}}_s\right] = \mathbb{E}\left[\int_0^T \varphi_s \hat{Z}_s \langle M\rangle^{\mathbb{H}}_s\right] = \mathbb{E}\left[\int_0^T \varphi_s \hat{Z}_s \langle M\rangle_s\right].$$

Finally, let us observe that the above equality is fulfilled for any $\mathbb{H}$-predictable process φ. Hence we can choose $\varphi = \hat{Z}$ and get

$$\mathbb{E}\left[\int_0^T |\hat{Z}_s|^2 \mathrm{d}\langle M\rangle_s\right] = \mathbb{E}\left[\int_0^T \hat{Z}_s \tilde{Z}_s \mathrm{d}\langle M\rangle_s\right].$$

Then, by Cauchy-Schwarz inequality we obtain

$$\mathbb{E}\left[\int_0^T |\hat{Z}_s|^2 \mathrm{d}\langle M\rangle_s\right] \leq \left\{\mathbb{E}\left[\int_0^T |\tilde{Z}_s|^2 \mathrm{d}\langle M\rangle_s\right]\right\}^{\frac{1}{2}} \left\{\mathbb{E}\left[\int_0^T |\hat{Z}_s|^2 \mathrm{d}\langle M\rangle_s\right]\right\}^{\frac{1}{2}}$$

which in turn implies

$$\mathbb{E}\left[\int_0^T |\hat{Z}_s|^2 \mathrm{d}\langle M\rangle_s\right] \leq \mathbb{E}\left[\int_0^T |\tilde{Z}_s|^2 \mathrm{d}\langle M\rangle_s\right] < \infty.$$

□

Remark 2.15. *As shown in Section 4 of [9], in some cases it is possible to compute explicitly the Radon-Nikodým derivative of $L^{\mathbb{H}}$ with respect to $\langle M\rangle^{\mathbb{H}}$ that characterizes the component Z of solution. For instance, if $\langle M\rangle$ is of the form*

$$\langle M\rangle_t = \int_0^t a_s \mathrm{d}G_s, \quad t \in [0,T]$$

for some $\mathbb{F}$-predictable process $a = (a_t)_{0\leq t\leq T}$ and an increasing deterministic function G, then

$$Z_t = \frac{{}^p(\tilde{Z}_t a_t)}{{}^p a_t}, \quad t \in [0,T],$$

where the notation ${}^pX$ refers to the $\mathbb{H}$-predictable projection of the process X. Another meaningful example is given by assuming $\langle M\rangle$ to be $\mathbb{H}$-predictable. In this case, we have

$$Z_t = {}^p\tilde{Z}_t, \quad t \in [0,T].$$

2.2. **Some special cases.** We are now in the position to provide an explicit characterization of the solution to the BSDE (2.2) under partial information in terms of the one related to the corresponding BSDE in the case of full information when the driver f does not depend on z.

Proposition 2.16. *Suppose that the driver f is independent of z and let $(\tilde{Y}, \tilde{Z}, \tilde{O}) \in S^2_{\mathcal{F}}(0,T) \times \mathcal{M}^2_{\mathcal{F}}(0,T) \times \mathcal{L}^2_{\mathcal{F}}(0,T)$ be a solution to the following BSDE under complete information*

$$\tilde{Y}_t = \xi + \int_t^T f(s, \tilde{Y}_{s-}) \mathrm{d}\langle M \rangle_s - \int_t^T \tilde{Z}_s \mathrm{d}M_s - (\tilde{O}_T - \tilde{O}_t), \quad 0 \leq t \leq T, \tag{2.14}$$

where $\tilde{O} = (\tilde{O}_t)_{0 \leq t \leq T}$ is a square-integrable $\mathbb{F}$-martingale strongly orthogonal to M. Set $L_t := \int_0^t \tilde{Z}_s \mathrm{d}\langle M \rangle_s$ for each $t \in [0,T]$. Then, the triplet

$$(Y, Z, O) = \left(\tilde{Y}, \frac{\mathrm{d}L^{\mathbb{H}}}{\mathrm{d}\langle M \rangle^{\mathbb{H}}}, \tilde{O} + B \right),$$

where $B = \int (\tilde{Z}_s - Z_s) \mathrm{d}M_s$ is a square-integrable $\mathbb{F}$-martingale weakly orthogonal to M, is a solution to the BSDE

$$Y_t = \xi + \int_t^T f(s, Y_{s-}) \mathrm{d}\langle M \rangle_s - \int_t^T Z_s \mathrm{d}M_s - (O_T - O_t), \quad 0 \leq t \leq T, \tag{2.15}$$

under partial information in the sense of Definition 2.3.

Proof. It is a direct consequence of Proposition 2.14. □

We conclude this subsection by applying Proposition 2.14 to provide existence of the solution to a BSDE under partial information in the special case where $\mathcal{H}_t = \mathcal{F}_{(t-\tau)^+}$ for each $t \in [0,T]$, with $\tau \in (0,T)$ being a fixed delay, the driver does not depend on y and $\langle M \rangle$ and $f(\cdot, \cdot, z)$ are $\mathbb{H}$-predictable processes. This approach allows us to weaken the assumptions required in Theorem 2.12. More precisely, we just require that f satisfies a sublinear growth condition in z.

Without loss of generality, we take $T = \tau N$, with $N \in \mathbb{N}$. We will solve backwardly equation (2.13) on each interval $I_j = [(j-1)\tau, j\tau]$, $j \in \{1, \dots, N\}$. To this aim we need a preliminary Lemma.

Lemma 2.17. *Let $\mathcal{H}_t = \mathcal{F}_{(t-\tau)^+}$, for each $t \in [0,T]$, with $\tau \in (0,T)$ being a fixed delay and assume that $\langle M \rangle_T \leq C(T)$ $\mathbb{P}$-a.s., where $C(T)$ is a positive constant depending on T. Let $\langle M \rangle$ and $f(\cdot, \cdot, z)$ be $\mathbb{H}$-predictable and f to satisfy a sublinear growth condition with respect to z uniformly in (ω, t), i.e.*

$\exists\, C \geq 0$ *such that* $\forall z \in \mathbb{R}$, $|f(\omega, t, z)|^2 \leq C(1 + |z|^2) \quad (\mathbb{P} \otimes \langle M \rangle) - a.e.$ *on* $\Omega \times [0,T]$;

Let $\xi^j \in L^2(\Omega, \mathcal{F}_{j\tau}, \mathbb{P}; \mathbb{R})$. Then there exists a solution $(\tilde{Y}^j, \tilde{Z}^j, \tilde{O}^j) \in \mathcal{S}^2_{\mathcal{F}}((j-1)\tau, j\tau) \times \mathcal{M}^2_{\mathcal{F}}((j-1)\tau, j\tau) \times \mathcal{L}^2_{\mathcal{F}}((j-1)\tau, j\tau)$ to the problem under complete information

$$\tilde{Y}^j_t = \xi^j + \int_t^{j\tau} f(s, {}^p\tilde{Z}^j_s) \mathrm{d}\langle M \rangle_s - \int_t^{j\tau} \tilde{Z}^j_s \mathrm{d}M_s - (\tilde{O}^j_{j\tau} - \tilde{O}^j_t), \quad (j-1)\tau \leq t \leq j\tau, \tag{2.16}$$

where $\tilde{O}^j$ is strongly orthogonal to M and ${}^p\tilde{Z}^j$ denotes the $\mathbb{H}$-predictable projection of $\tilde{Z}^j$, that is, ${}^p\tilde{Z}^j_t = \mathbb{E}\left[\tilde{Z}^j_t | \mathcal{H}_{t^-}\right]$, for every $t \in [0,T]$.

Proof. According to the Galtchouk-Kunita-Watanabe decomposition of ξ^j under full information, there exists $\tilde{Z}^j \in \mathcal{M}^2_{\mathcal{F}}((j-1)\tau, j\tau)$ such that

$$\xi^j = \mathbb{E}\left[\xi^j | \mathcal{F}_{(j-1)\tau}\right] + \int_{(j-1)\tau}^{j\tau} \tilde{Z}^j_s \mathrm{d}M_s + \left(\tilde{O}^j_{j\tau} - \tilde{O}^j_{(j-1)\tau}\right), \tag{2.17}$$

where $\tilde{O}^j \in L^2_{\mathcal{F}}((j-1)\tau, j\tau)$ is strongly orthogonal to M. For every $t \in [(j-1)\tau, j\tau]$, we set

$$Y^j_t = \mathbb{E}\left[\xi^j | \mathcal{F}_t\right] + \int_t^{j\tau} f(s, {}^p\tilde{Z}^j_s)\mathrm{d}\langle M\rangle_s. \tag{2.18}$$

Let us observe that $Y^j \in S^2_{\mathcal{F}}((j-1)\tau, j\tau)$. In fact, since $\int_t^{j\tau} f(s, {}^p\tilde{Z}^j_s)\mathrm{d}\langle M\rangle_s$ is $\mathcal{F}_{(j-1)\tau}$-measurable Y^j turns out to be $\mathbb{F}$-adapted. By the sublinear growth condition on f, Jensen's inequality and the property of the $\mathbb{H}$-predictable projection, we get

$$\begin{aligned}
\mathbb{E}\left[\int_{(j-1)\tau}^{j\tau} |f(s, {}^p\tilde{Z}^j_s)|^2 \mathrm{d}\langle M\rangle_s\right] &\leq \mathbb{E}\left[\int_t^{j\tau} C(1 + |{}^p\tilde{Z}^j_s|^2)\mathrm{d}\langle M\rangle_s\right] \\
&\leq \mathbb{E}\left[\int_{(j-1)\tau}^{j\tau} C(1 + {}^p(|\tilde{Z}^j|^2_s))\mathrm{d}\langle M\rangle_s\right] \\
&= C\,\mathbb{E}\left[\langle M\rangle_{j\tau} - \langle M\rangle_{(j-1)\tau} + \int_{(j-1)\tau}^{j\tau} |\tilde{Z}^j|^2_s \mathrm{d}\langle M\rangle_s\right] < \infty,
\end{aligned}$$

and by performing the same computation as in the proof of Lemma 2.10, we finally obtain

$$\mathbb{E}\left[\sup_{(j-1)\tau \leq t \leq j\tau} |Y^j_t|^2\right] \leq 8\mathbb{E}\left[|\xi^j|^2\right] + 8C(T)\mathbb{E}\left[\int_{(j-1)\tau}^{j\tau} |f(s, {}^p\tilde{Z}^j_s)|^2 \mathrm{d}\langle M\rangle_s\right] < \infty.$$

We now take the conditional expectation with respect to $\mathcal{F}_t$ in (2.17) and for each $t \in [(j-1)\tau, j\tau]$ we obtain

$$\mathbb{E}\left[\xi^j | \mathcal{F}_t\right] - \mathbb{E}\left[\xi^j | \mathcal{F}_{(j-1)\tau}\right] = \int_{(j-1)\tau}^{t} \tilde{Z}^j_s \mathrm{d}M_s + \left(\tilde{O}^j_t - \tilde{O}^j_{(j-1)\tau}\right). \tag{2.19}$$

At this stage, subtracting (2.19) and (2.17) yields

$$\mathbb{E}\left[\xi^j | \mathcal{F}_t\right] - \xi^j = -\int_{(j-1)\tau}^{t} \tilde{Z}^j_s \mathrm{d}M_s - \left(\tilde{O}^j_{j\tau t} - \tilde{O}^j_t\right)$$

and using (2.18) we get

$$Y^j_t - \int_t^{j\tau} f(s, {}^p\tilde{Z}^j_s)\mathrm{d}\langle M\rangle_s - \xi^j = -\int_{(j-1)\tau}^{t} \tilde{Z}^j_s \mathrm{d}M_s - \left(\tilde{O}^j_{j\tau} - \tilde{O}^j_t\right),$$

which concludes the proof. □

We are now in the position to state the following result.

Proposition 2.18. *Let $\mathcal{H}_t = \mathcal{F}_{(t-\tau)^+}$, for each $t \in [0,T]$, with $\tau \in (0,T)$ being a fixed delay and assume that $\langle M\rangle_T \leq C(T)$ $\mathbb{P}$-a.s., where $C(T)$ is a positive constant depending on T. Let $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$, $\langle M\rangle$ and $f(\cdot,\cdot,z)$ be $\mathbb{H}$-predictable and f to satisfy a sublinear growth condition with respect to z uniformly in (ω, t), i.e.*

$\exists\, C \geq 0$ *such that* $\forall z \in \mathbb{R}$, $|f(\omega,t,z)|^2 \leq C(1+|z|^2)$ $(\mathbb{P}\otimes\langle M\rangle)-a.e.$ *on* $\Omega\times[0,T]$.

Then, there exists a solution $(Y,Z,O)\in S^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{H}}(0,T)\times\mathcal{L}^2_{\mathcal{F}}(0,T)$ *to the BSDE under restricted information*

$$Y_t=\xi+\int_t^T f(s,Z_s)\mathrm{d}\langle M\rangle_s-\int_t^T Z_s\mathrm{d}M_s-(\tilde{O}_T-\tilde{O}_t). \tag{2.20}$$

Proof. We apply Lemma 2.17. Set $\xi^N=\xi$, and $\xi^j=\tilde{Y}^{j+1}_{j\tau}$, $j=1,2...N-1$, where $(\tilde{Y}^j,\tilde{Z}^j,\tilde{O}^j)\in\mathcal{S}^2_{\mathcal{F}}((j-1)\tau,j\tau)\times\mathcal{M}^2_{\mathcal{F}}((j-1)\tau,j\tau)\times\mathcal{L}^2_{\mathcal{F}}((j-1)\tau,j\tau)$ is the solution of the problem under complete information (2.16).

Set $\tilde{Y}_t:=\sum_{j=1}^N Y^j_t\mathbf{1}_{\{t\in[(j-1)\tau,j\tau)\}}$, $\tilde{Z}_t:=\sum_{j=1}^N Y^j_t\mathbf{1}_{\{t\in((j-1)\tau,j\tau]\}}$, $\tilde{O}_t:=\sum_{j=1}^N O^j_t\mathbf{1}_{\{t\in[(j-1)\tau,j\tau)\}}$. Then, we get that the triplet

$$(\tilde{Y},\tilde{Z},\tilde{O})\in\mathcal{S}^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{F}}(0,T)\times\mathcal{L}^2_{\mathcal{F}}(0,T)$$

is a solution to the problem under complete information

$$\tilde{Y}_t=\xi+\int_t^T f(s,{}^p\tilde{Z}_s)\mathrm{d}\langle M\rangle_s-\int_t^T\tilde{Z}_s\mathrm{d}M_s-(\tilde{O}_T-\tilde{O}_t),\quad 0\leq t\leq T. \tag{2.21}$$

Finally by applying Proposition 2.14, the triplet $(Y,Z,O)=(\tilde{Y},{}^p\tilde{Z},\tilde{O}+B)\in\mathcal{S}^2_{\mathcal{F}}(0,T)\times\mathcal{M}^2_{\mathcal{H}}(0,T)\times\mathcal{L}^2_{\mathcal{F}}(0,T)$, where $B=\int(\tilde{Z}_s-Z_s)\mathrm{d}M_s$ is a square-integrable $\mathbb{F}$-martingale weakly orthogonal to M, solves the BSDE (2.20) under restricted information. □

In the next section, we will apply the existence and uniqueness results obtained for BSDEs to derive the Föllmer-Schweizer decomposition in a partial information framework and discuss a financial application. More precisely, we will study the hedging problem of a contingent claim in incomplete markets when the underlying price process is given by a general $\mathbb{F}$-semimartingale and there are restrictions on the available information to traders.

3. Local Risk-Minimization under restricted information

Let us fix a probability space $(\Omega,\mathcal{F},\mathbb{P})$ endowed with a filtration $\mathbb{F}:=(\mathcal{F}_t)_{0\leq t\leq T}$ satisfying the usual conditions of right-continuity and completeness. Here $T>0$ denotes a fixed and finite time horizon; furthermore, we assume that $\mathcal{F}=\mathcal{F}_T$. We consider a financial market with one riskless asset with (discounted) price 1 and a risky asset whose (discounted) price S is described by an $\mathbb{R}$-valued square-integrable (càdlàg) $\mathbb{F}$-semimartingale $S=(S_t)_{0\leq t\leq T}$ satisfying the so-called structure condition (SC), that is

$$S_t=S_0+M_t+\int_0^t\alpha_s\mathrm{d}\langle M\rangle_s,\quad 0\leq t\leq T, \tag{3.1}$$

where $M=(M_t)_{0\leq t\leq T}$ is an $\mathbb{R}$-valued square-integrable (càdlàg) $\mathbb{F}$-martingale with $M_0=0$ and $\mathbb{F}$-predictable quadratic variation process denoted by $\langle M\rangle=(\langle M,M\rangle)_{0\leq t\leq T}$ and $\alpha=(\alpha_t)_{0\leq t\leq T}$ is an $\mathbb{F}$-predictable process such that $\mathbb{E}\left[\int_0^T|\alpha_t|^2\mathrm{d}\langle M\rangle_t\right]<\infty$.

Remark 3.1. *It is known that the existence of an equivalent martingale measure for the risky asset price process S implies that S is an $\mathbb{F}$-semimartingale under the basic measure $\mathbb{P}$. Then, the semimartingale structure for S is a natural assumption in a financial market model which ensures the absence of arbitrage opportunities. If in addition, S has continuous trajectories or has càdlàg paths and the following condition holds*

$$\mathbb{E}\left[\sup_{t\in[0,T]} S_t^2\right] < \infty,$$

then S satisfies the structure condition (SC), see page 24 of [1] and Theorem 1 in [16].

In this framework we consider a contingent claim whose payoff is represented by a random variable $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$. Under the condition that the mean-variance tradeoff process $K = (K_t)_{0\leq t\leq T}$ defined by

$$K_t := \int_0^t \alpha_s^2 \mathrm{d}\langle M\rangle_s, \quad \forall t \in [0,T],$$

is uniformly bounded in t and ω, in Theorem 3.4 of [16] it is proved that every $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$ admits a strong Föllmer-Schweizer decomposition with respect to S, that is

$$\xi = \tilde{U}_0 + \int_0^T \beta_t \mathrm{d}S_t + \tilde{A}_T, \quad \mathbb{P}-\text{a.s.}, \tag{3.2}$$

where $\tilde{U}_0 \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; \mathbb{R})$, $\beta = (\beta_t)_{0\leq t\leq T}$ is an $\mathbb{F}$-predictable process such that the stochastic integral $\int \beta_t \mathrm{d}S_t$ is well-defined and it is a square-integrable $\mathbb{F}$-semimartingale and $\tilde{A} \in \mathcal{L}^2_{\mathcal{F}}(0,T)$ is strongly orthogonal to M, see (3.1). Moreover, it is known that every $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$ admits a decomposition (3.2) if and only if there exists a locally risk-minimizing hedging strategy (see e.g. [12, 23]) and in addition this decomposition plays an essential role in the variance-minimizing strategy computation (see [21] for further details).

Suppose now that the hedger does not have at her/his disposal the full information represented by $\mathbb{F}$; her/his strategy must be constructed from less information. This leads to a partial information framework. To describe this mathematically, we introduce an additional filtration $\mathbb{H} := (\mathcal{H}_t)_{0\leq t\leq T}$ satisfying the usual conditions and such that $\mathcal{H}_t \subseteq \mathcal{F}_t$, for every $t \in [0,T]$. Thanks to Theorem 2.12, we are now in the position to derive a similar decomposition in a partial information setting. We need the following additional hypothesis.

Assumption 3.2. *There exists a constant $\bar{K} \geq 0$ such that the process α in (3.1) satisfies:*

$$|\alpha_t(\omega)| \leq \bar{K}, \quad (\mathbb{P}\otimes\langle M\rangle) - a.e. \text{ on } \Omega\times[0,T].$$

Proposition 3.3. *Let Assumptions 2.8 and 3.2 hold. Then, every $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$ admits the following decomposition*

$$\xi = \bar{U}_0 + \int_0^T \beta_t^{\mathcal{H}} \mathrm{d}S_t + A_T, \quad \mathbb{P}-a.s., \tag{3.3}$$

where $\bar{U}_0 \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; \mathbb{R})$, $\beta^{\mathcal{H}} = (\beta_t^{\mathcal{H}})_{0\le t\le T} \in \mathcal{M}^2_{\mathcal{H}}(0,T)$ *and* $A \in \mathcal{L}^2_{\mathcal{F}}(0,T)$ *is weakly orthogonal to* M*.*

In the martingale case where $\alpha \equiv 0$ in (3.1), representation (3.3) corresponds to the Galtchouk-Kunita-Watanabe decomposition (2.4) of ξ under partial information. In the general semimartingale case, (3.3) is referred as the Föllmer-Schweizer decomposition of ξ with respect to S under partial information.

Proof. Let us consider the driver of the BSDE (2.2) under partial information given by $f(t,y,z) = -z\alpha$, where α is the bounded process introduced in (3.1). Since Assumption 2.1 is fulfilled, by Theorem 2.12 there exists a unique triplet (Y, Z, O) which solves the equation

$$Y_t = \xi - \int_t^T Z_s \alpha_s \mathrm{d}\langle M\rangle_s - \int_t^T Z_s \mathrm{d}M_s - (O_T - O_t), \quad 0 \le t \le T, \tag{3.4}$$

under partial information in the sense of Definition 2.3. Hence

$$\xi = Y_T = Y_0 + \int_0^T Z_s \alpha_s \mathrm{d}\langle M\rangle_s + \int_0^T Z_s \mathrm{d}M_s + O_T = Y_0 + \int_0^T Z_s \mathrm{d}S_s + O_T$$

and we obtain decomposition (3.3) by setting $\bar{U}_0 = Y_0$, $\beta_t^{\mathcal{H}} = Z_t$ and $A_t = O_t$, for every $t \in [0,T]$. □

Remark 3.4. *Note that if* Y *represents the wealth that satisfies the replication constraint* $Y_T = \xi$ $\mathbb{P}$*-a.s., the triplet* $(Y, \beta^{\mathcal{H}}, A)$ *may be interpreted as the* nonadjusted hedging strategy *against* ξ*. Clearly, the self-financing condition of the strategy is no longer ensured due to the presence of the cost* A*, see [11] for further details.*

We now study the relationship between the Föllmer-Schweizer decomposition of a contingent claim $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$ under partial information and the existence of a *locally risk-minimizing* strategy in a partial information framework. In the sequel, we will suppose that Assumptions 2.8 and 3.2 are in force.
In this setting, the amount $\theta = (\theta_t)_{0\le t\le T}$ invested by the agent in the risky asset has to be adapted to the information flow $\mathbb{H}$ and such that the stochastic integral $\int \theta_u \mathrm{d}S_u$ turns out to be a square-integrable $\mathbb{F}$-semimartingale. By Assumption 2.8 and boundedness of α, we will look at the class of processes θ such that $\theta \in \mathcal{M}^2_{\mathcal{H}}(0,T)$. Indeed,

$$\begin{aligned}
\mathbb{E}\left[\int_0^T \theta_s^2 \mathrm{d}\langle M\rangle_s + \left(\int_0^T |\theta_s||\alpha_s| \mathrm{d}\langle M\rangle_s\right)^2\right] &\le \mathbb{E}\left[\int_0^T \theta_s^2 \mathrm{d}\langle M\rangle_s + \bar{K}^2 \left(\int_0^T |\theta_s| \mathrm{d}\langle M\rangle_s\right)^2\right] \\
&\le \mathbb{E}\left[\int_0^T \theta_s^2 \mathrm{d}\langle M\rangle_s + \bar{K}^2 \int_0^T \theta_s^2 \mathrm{d}\langle M\rangle_s \cdot \langle M\rangle_T\right] \\
&\le \left(1 + \bar{K}^2 \rho(T)\right) \mathbb{E}\left[\int_0^T \theta_s^2 \mathrm{d}\langle M\rangle_s\right].
\end{aligned}$$

Clearly, in this case Assumption 2.8 can be weakened by requiring that $\langle M\rangle_T \le C(T)$ $\mathbb{P}$-a.s., for a positive constant $C(T)$ depending on T.

Definition 3.5. *An* $(\mathbb{H},\mathbb{F})$-admissible strategy *is a pair* $\Psi=(\theta,\eta)$ *where* $\theta\in\mathcal{M}^2_{\mathcal{H}}(0,T)$ *and* $\eta=(\eta_t)_{0\le t\le T}$ *is a real-valued* $\mathbb{F}$*-adapted process such that the value process* $V(\Psi):=\theta S+\eta$ *is right-continuous and satisfies* $V_t(\Psi)\in L^2(\Omega,\mathcal{F}_t,\mathbb{P};\mathbb{R})$ *for each* $t\in[0,T]$.

Remark 3.6. *We assume that the agent has at her/his disposal the information flow* $\mathbb{H}$ *about trading in the risky asset while a complete information about trading in the riskless asset.*

Given an $(\mathbb{H},\mathbb{F})$-admissible strategy Ψ, the associated *cost process* $C(\Psi)=(C_t(\Psi))_{0\le t\le T}$ is defined by

$$C_t(\Psi)=V_t(\Psi)-\int_0^t\theta_s\mathrm{d}S_s,\quad\forall t\in[0,T].$$

Here $C_t(\Psi)$ describes the total costs incurred by Ψ over the interval $[0,t]$. The $\mathbb{H}$-*risk process* $R^{\mathcal{H}}(\Psi)=(R^{\mathcal{H}}_t(\Psi))_{0\le t\le T}$ of Ψ is then defined by

$$R^{\mathcal{H}}_t(\Psi):=\mathbb{E}\left[\left(C_T(\Psi)-C_t(\Psi)\right)^2\Big|\,\mathcal{H}_t\right],\quad\forall t\in[0,T].\tag{3.5}$$

Although $(\mathbb{H},\mathbb{F})$-admissible strategies Ψ with $V_T(\Psi)=\xi$ will in general not be self-financing, it turns out that good $(\mathbb{H},\mathbb{F})$-admissible strategies are still self-financing on average in the following sense.

Definition 3.7. *An* $(\mathbb{H},\mathbb{F})$*-admissible strategy* Ψ *is called* mean-self-financing *if its cost process* $C(\Psi)$ *is an* $\mathbb{F}$*-martingale.*

Inspired by [20], an $(\mathbb{H},\mathbb{F})$-admissible strategy Ψ is called $(\mathbb{H},\mathbb{F})$-*locally risk-minimizing* if, for any $t<T$, the remaining risk $R^{\mathcal{H}}(\Psi)$, see (3.5), is minimal under all infinitesimal perturbations of the strategy at time t. For further details, we refer to Definition A.2 in Appendix.

Proposition 3.8. *Suppose that* $\langle M\rangle$ *is* $\mathbb{P}$*-a.s. strictly increasing. Let* $\xi\in L^2(\Omega,\mathcal{F}_T,\mathbb{P};\mathbb{R})$ *be a contingent claim and* Ψ *an* $(\mathbb{H},\mathbb{F})$*-admissible strategy with* $V_T(\Psi)=\xi$ $\mathbb{P}$*-a.s.. Then* Ψ *is* $(\mathbb{H},\mathbb{F})$*-locally risk-minimizing if and only if* Ψ *is mean-self-financing and the* $\mathbb{F}$*-martingale* $C(\Psi)$ *is weakly orthogonal to* M.

Proof. For the proof, we refer to Section A in Appendix. □

The previous result motivates the following.

Definition 3.9. *Let* $\xi\in L^2(\Omega,\mathcal{F}_T,\mathbb{P};\mathbb{R})$ *be a contingent claim. An* $(\mathbb{H},\mathbb{F})$*-admissible strategy* Ψ *with* $V_T(\psi)=\xi$ $\mathbb{P}$*-a.s. is called* $(\mathbb{H},\mathbb{F})$-optimal *for* ξ *if* Ψ *is mean-self-financing and the* $\mathbb{F}$*-martingale* $C(\Psi)$ *is weakly orthogonal to* M.

The next result ensures that the existence of an $(\mathbb{H},\mathbb{F})$-*optimal* strategy is equivalent to the decomposition (3.3) of the contingent claim ξ. In the case of full information, an analogous result can be found in [12].

Proposition 3.10. *A contingent claim $\xi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}; \mathbb{R})$ admits an $(\mathbb{H}, \mathbb{F})$-optimal strategy $\Psi = (\theta, \eta)$ with $V_T(\Psi) = \xi$ $\mathbb{P}$-a.s. if and only if ξ can be written as*

$$\xi = U_0 + \int_0^T \beta_t^{\mathcal{H}} \mathrm{d}S_t + A_T, \quad \mathbb{P} - a.s., \tag{3.6}$$

with $U_0 \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; \mathbb{R})$, $\beta^{\mathcal{H}} \in \mathcal{M}^2_{\mathcal{H}}(0,T)$ and $A \in \mathcal{L}^2_{\mathcal{F}}(0,T)$ weakly orthogonal to M. The strategy Ψ is then given by

$$\theta_t = \beta_t^{\mathcal{H}}, \quad 0 \leq t \leq T$$

with minimal cost

$$C_t(\Psi) = U_0 + A_t, \quad 0 \leq t \leq T.$$

If (3.6) *holds, the optimal portfolio value is*

$$V_t(\Psi) = C_t(\Psi) + \int_0^t \theta_s \mathrm{d}S_s = U_0 + \int_0^t \beta_s^{\mathcal{H}} \mathrm{d}S_s + A_t, \quad 0 \leq t \leq T$$

and

$$\eta_t = V_t(\Psi) - \beta_t^{\mathcal{H}} S_t, \quad 0 \leq t \leq T.$$

Proof. Suppose that Ψ is an $(\mathbb{H}, \mathbb{F})$-optimal strategy with $V_T(\Psi) = \xi$ $\mathbb{P}$-a.s.. Then, the replication constraint yields

$$\xi = V_T(\Psi) = C_T(\Psi) + \int_0^T \theta_s \mathrm{d}S_s = C_0(\Psi) + \int_0^T \theta_s \mathrm{d}S_s + \left(C_T(\Psi) - C_0(\Psi)\right), \quad \mathbb{P} - \text{a.s.}. \tag{3.7}$$

Since Ψ is an $(\mathbb{H}, \mathbb{F})$-optimal strategy, by Proposition 3.8 we know that the process $C(\Psi) - C_0(\Psi)$ is a square-integrable $\mathbb{F}$-martingale weakly orthogonal to M that is in addition null at zero. Hence (3.7) is indeed the Föllmer-Schweizer decomposition of ξ with respect to S under partial information with $\beta^{\mathcal{H}} = \theta$ and $A = C(\Psi) - C_0(\Psi)$.
We now assume that (3.6) holds. Then, we choose

$$\begin{aligned} &\theta_t = \beta_t^{\mathcal{H}}, \quad t \in [0,T], \\ &\eta_t = U_0 + A_t - \beta_t^{\mathcal{H}} S_t - \int_0^t \beta_s^{\mathcal{H}} \mathrm{d}S_s, \quad t \in [0,T]. \end{aligned}$$

Thus, the strategy $\Psi = (\beta^{\mathcal{H}}, \eta)$ is such that the associated cost is given by

$$C_t(\Psi) = V_t(\Psi) - \int_0^t \beta_s^{\mathcal{H}} \mathrm{d}S_s = U_0 + A_t,$$

for every $t \in [0,T]$. In particular, $C_T(\Psi) = U_0 + A_T$. Hence $C(\Psi)$ is an $\mathbb{F}$-martingale weakly orthogonal to M and this implies that Ψ is an $(\mathbb{H}, \mathbb{F})$-optimal strategy. □

Remark 3.11. *As a consequence of Proposition 3.10 and Theorem 2.12, under Assumptions 2.8 and 3.2, we can characterize, the $(\mathbb{H}, \mathbb{F})$-optimal strategy $\Psi = (\theta, \eta)$, the optimal portfolio value $V(\Psi)$ and the corresponding minimal cost $C(\Psi)$, in terms of the unique solution (Y, Z, O) to the BSDE* (2.2) *with the particular choice of $f(t, y, z) = -\alpha_t z$; more precisely, $V(\Psi) = Y$, $\theta = Z$ and $C(\Psi) = O + Y_0$.*

By applying Proposition 2.14 (with the particular choice of $f(t,y,z) = -\alpha_t z$) the $(\mathbb{H},\mathbb{F})$-optimal strategy may be expressed in terms of the solution of a problem under full information.

Proposition 3.12. *Let Assumptions 2.8 and 3.2 hold. Let* $(\tilde{Y},\tilde{Z},\tilde{O}) \in \mathcal{S}^2_{\mathcal{F}}(0,T) \times \mathcal{M}^2_{\mathcal{F}}(0,T) \times \mathcal{L}^2_{\mathcal{F}}(0,T)$ *be a solution to the problem under complete information*

$$\tilde{Y}_t = \xi - \int_t^T \hat{Z}_s \alpha_s \mathrm{d}\langle M\rangle_s - \int_t^T \tilde{Z}_s \mathrm{d}M_s - (\tilde{O}_T - \tilde{O}_t), \quad 0 \le t \le T, \tag{3.8}$$

where $\tilde{O}$ *is strongly orthogonal to* M *and*

$$\hat{Z}_t := \frac{\mathrm{d}L^{\mathbb{H}}_t}{\mathrm{d}\langle M\rangle^{\mathbb{H}}_t} \quad L_t := \int_0^t \tilde{Z}_s \mathrm{d}\langle M\rangle_s, \quad 0 \le t \le T.$$

Then the $(\mathbb{H},\mathbb{F})$*-optimal strategy* $\Psi = (\beta^{\mathcal{H}},\eta)$*, the optimal portfolio value and the minimal cost are given by*

$$\beta^{\mathcal{H}}_t = \hat{Z}_t, \quad V_t(\Psi) = \tilde{Y}_t, \quad C_t(\Psi) = \tilde{Y}_0 + \tilde{O}_t + \int_0^t (\tilde{Z}_s - \hat{Z}_s)\mathrm{d}M_s \quad \forall t \in [0,T],$$

respectively.

Proof. By Proposition 2.14 we get the the triplet

$$(Y,Z,O) = \left(\tilde{Y},\hat{Z},\tilde{O} + B\right),$$

where $B = \int(\tilde{Z}_s - \hat{Z}_s)\mathrm{d}M_s$ is a square-integrable $\mathbb{F}$-martingale weakly orthogonal to M, is a solution to the BSDE (2.2) under partial information with the particular choice of $f(t,y,z) = -\alpha_t z$.

Finally, by uniqueness of the solution to this equation and Remark 3.11 the thesis follows. □

Remark 3.13. *Let us observe that the process* $\tilde{Z}$ *coincides with the optimal strategy under full information,* β*, only in the particular case where* S *is an* $\mathbb{F}$*-martingale, i.e.* $S = M$ *(see [9]). In fact, in the semimartingale case,* β *is given by the second component of the solution to the BSDE under full information with the choice* $f(t,y,z) = -\alpha_t z$ *which differs from equation (3.8) that is not a BSDE.*

3.1. **Local risk-minimization under complete information.** Under full information and in the case where the stock price process S has continuous trajectories, the locally risk-minimizing strategy can be computed via the Galtchouk-Kunita-Watanabe decompositon of the contingent claim with respect to the minimal martingale measure (in short MMM) $\mathbb{P}^*$, see e.g. Theorem 3.5 of [23]. This is a consequence of the fact that the MMM preserves orthogonality, which means that any $(\mathbb{P},\mathbb{F})$-martingale strongly orthogonal to the martingale part of S under $\mathbb{P}$ turns out to be a $(\mathbb{P}^*,\mathbb{F})$-martingale strongly orthogonal to S under $\mathbb{P}^*$. We emphasize that this is no longer true in general if S has jumps. However, we are able to characterize the optimal portfolio value in terms of the MMM for S even in presence of jumps.

Let us recall the definition of the MMM.

Definition 3.14. *An equivalent martingale measure $\mathbb{P}^*$ for S with square-integrable density $\mathrm{d}\mathbb{P}^*/\mathrm{d}\mathbb{P}$ is called* minimal martingale measure *(for S) if $\mathbb{P}^* = \mathbb{P}$ on $\mathcal{F}_0$ and if every $(\mathbb{P},\mathbb{F})$-martingale $\widetilde{A}$ which is square-integrable and strongly orthogonal to the martingale part of S is also a $(\mathbb{P}^*,\mathbb{F})$-martingale. We call $\mathbb{P}^*$* orthogonality-preserving *if $\widetilde{A}$ is also strongly orthogonal to S under $\mathbb{P}^*$.*

From now on we assume an additional condition on the jump sizes of the martingale part M of S which ensures the existence of the MMM for S. More precisely, we make the following assumption:

$$1 - \alpha_t \Delta M_t > 0 \quad \mathbb{P} - \text{a.s.} \quad \forall t \in [0,T]. \tag{3.9}$$

Hence by the Ansel-Strickel Theorem, see [1], there exists the minimal martingale measure $\mathbb{P}^*$ for S defined by

$$\frac{\mathrm{d}\mathbb{P}^*}{\mathrm{d}\mathbb{P}}\Big|_{\mathcal{F}_t} = \tilde{L}_t := \mathcal{E}\left(-\int \alpha_r \mathrm{d}M_r\right)_t, \quad t \in [0,T], \tag{3.10}$$

where $\mathcal{E}$ denotes the Doléans-Dade exponential. Let us observe that by Assumptions 2.8 and 3.2 the following estimate holds

$$\mathbb{E}\left[e^{\int_0^T \alpha_t^2 \mathrm{d}\langle M\rangle_t}\right] = e^{\bar{K}\rho(T)}$$

which implies that the nonnegative $(\mathbb{P},\mathbb{F})$-local martingale $\tilde{L}$ is in fact a square-integrable $(\mathbb{P},\mathbb{F})$-martingale, see e.g. [18].

Proposition 3.15. *Let Assumptions 2.8, 3.2 and equation (3.9) hold, $\xi \in L^2(\Omega,\mathcal{F}_T,\mathbb{P};\mathbb{R})$, $\mathbb{H} = \mathbb{F}$ and assume the $\langle M\rangle$ to be $\mathbb{P}$-a.s. strictly increasing. Then there exists the (classical) locally risk-minimizing strategy $\Psi = (\theta,\eta)$ for ξ and the optimal portfolio value $V^{\mathcal{F}}(\Psi)$ can be computed via the MMM as*

$$V_t^{\mathcal{F}}(\Psi) = E^{\mathbb{P}^*}[\xi|\mathcal{F}_t] \quad \forall t \in [0,T],$$

where the notation $E^{\mathbb{P}^*}[\cdot|\mathcal{F}_t]$ denotes the conditional expectation with respect to $\mathcal{F}_t$ computed under $\mathbb{P}^*$.

Proof. First let us observe that $\xi \in L^2(\Omega,\mathcal{F}_T,\mathbb{P};\mathbb{R})$ and $\tilde{L}$ square-integrable $(\mathbb{P},\mathbb{F})$-martingale imply that $\xi \in L^1(\Omega,\mathcal{F}_T,\mathbb{P}^*;\mathbb{R})$.
By Propositions 3.3, 3.8 and 3.10 we deduce the existence of the (classical) locally risk-minimizing strategy $\Psi = (\theta,\eta)$. Consider the Föllmer-Schweizer decomposition of ξ under full information:

$$\xi = \widetilde{U}_0 + \int_0^T \beta_t \mathrm{d}S_t + \widetilde{A}_T, \quad \mathbb{P} - \text{a.s.}, \tag{3.11}$$

where $\widetilde{U}_0 \in L^2(\Omega,\mathcal{F}_0,\mathbb{P};\mathbb{R})$, β is an $\mathbb{F}$-predictable process such that $\mathbb{E}\left[\int_0^T \beta_s^2 \mathrm{d}\langle M\rangle_s\right] < \infty$ and $\widetilde{A} \in \mathcal{L}^2_{\mathcal{F}}(0,T)$ is strongly orthogonal to M. Then, $\theta = \beta$ in (3.11) and the optimal portfolio value $V^{\mathcal{F}}(\Psi)$ satisfies for each $t \in [0,T]$

$$V_t^{\mathcal{F}}(\Psi) = \widetilde{U}_0 + \int_0^t \beta_u \mathrm{d}S_u + \widetilde{A}_t, \tag{3.12}$$

with $\widetilde{A} \in \mathcal{L}^2_{\mathcal{F}}(0,T)$ strongly orthogonal to M. Since $\int \beta_r \mathrm{d}M_r$ and $\tilde{L}$ are $(\mathbb{P},\mathbb{F})$-square integrable martingales, then $\int \beta_r \mathrm{d}S_r$ is a $(\mathbb{P}^*,\mathbb{F})$-martingale (see the proof of Theorem 3.14 in [12]). Therefore, the definition of MMM yields that the optimal portfolio value $V^{\mathcal{F}}(\Psi)$ turns out to be a $(\mathbb{P}^*,\mathbb{F})$-martingale and as a consequence, we get

$$V_t^{\mathcal{F}}(\Psi) = E^{\mathbb{P}^*}[V_T^{\mathcal{F}}(\Psi)|\mathcal{F}_t] = E^{\mathbb{P}^*}[\xi|\mathcal{F}_t], \quad t \in [0,T].$$

□

Remark 3.16. *Let us observe that such a result cannot be extended to the partial information framework, since in the Föllmer-Schweizer decomposition of ξ under partial information (see equation (3.3)) the $\mathbb{F}$-martingale A is only weakly orthogonal to M and so A is not in general a $(\mathbb{P}^*,\mathbb{F})$-martingale.*

Remark 3.17. *Proposition 3.15 may be useful to compute the locally risk-minimizing strategy under full information, since by (3.12), it may be expressed using the predictable covariation under $\mathbb{P}$ of $V^{\mathcal{F}}(\Psi)$ and S, i.e.*

$$\beta_t = \frac{\mathrm{d}\langle V^{\mathcal{F}}(\Psi), S\rangle^{\mathbb{P}}}{\mathrm{d}\langle S\rangle^{\mathbb{P}}}, \quad t \in [0,T].$$

See [24] and references therein for explicit solutions in exponential Lévy models.

Appendix A. Technical Results

Here we clarify the concept of an $(\mathbb{H},\mathbb{F})$-locally risk-minimizing strategy. As the original version given in the case full information, see e.g. [20], this concept translates the idea that changing an optimal strategy over a small time interval should lead to an increase of risk, at least asymptotically.

Definition A.1. *A* small perturbation *is an $(\mathbb{H},\mathbb{F})$-admissible strategy $\Delta = (\delta,\gamma)$ such that δ is bounded, the variation of $\int \delta_u \alpha_u \mathrm{d}\langle M\rangle_u$ is bounded (uniformly in t and ω) and $\delta_T = \gamma_T = 0$. For any subinterval $(s,t]$ of $[0,T]$, we then define the small perturbation*

$$\Delta|_{(s,t]} := \left(\delta \mathbf{1}_{(s,t]}, \gamma \mathbf{1}_{[s,t)}\right).$$

To explain the notion of a local variation of an $(\mathbb{H},\mathbb{F})$-admissible strategy, we consider partitions $\tau = (t_i)_{0\le i\le N}$ of the interval $[0,T]$. Such partitions will always satisfy

$$0 = t_0 < t_1 < \ldots < t_N = T.$$

Definition A.2. *For an $(\mathbb{H},\mathbb{F})$-admissible strategy Ψ, a small perturbation Δ and a partition τ of $[0,T]$, we set*

$$r^{\tau}_{\mathcal{H}}(\Psi,\Delta) := \sum_{t_i,t_{i+1}\in\tau} \frac{R^{\mathcal{H}}_{t_i}\left(\Psi + \Delta|_{(t_i,t_{i+1}]}\right) - R^{\mathcal{H}}_{t_i}(\Psi)}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}} - \langle M\rangle_{t_i} \middle|\, \mathcal{H}_{t_i}\right]} \mathbf{1}_{(t_i,t_{i+1}]}. \tag{A.1}$$

The strategy Ψ is called $(\mathbb{H},\mathbb{F})$-locally risk-minimizing if

$$\liminf_{n\to\infty} r^{\tau_n}_{\mathcal{H}}(\Psi,\Delta) \ge 0, \quad (\mathbb{P}\otimes\langle M\rangle) - \text{a.e. on } \Omega\times[0,T]$$

for every small perturbation Δ and every increasing sequence $(\tau_n)_{n\in\mathbb{N}}$ of partitions of $[0,T]$ tending to identity.

Remark A.3. *If an $(\mathbb{H},\mathbb{F})$-admissible strategy $\Psi=(\theta,\eta)$ is mean-self-financing, that is $C(\Psi)$ is an $\mathbb{F}$-martingale, Ψ is uniquely determined by θ. Indeed, since by the replication constraint we have*

$$C_T(\Psi)=V_T(\Psi)-\int_0^T\theta_s\mathrm{d}S_s=\xi-\int_0^T\theta_s\mathrm{d}S_s,$$

then, by the mean-self-financing property, we get

$$C_t(\Psi)=\mathbb{E}\left[\xi-\int_0^T\theta_s\mathrm{d}S_s\middle|\mathcal{F}_t\right],\quad t\in[0,T].$$

Hence we can write $C(\theta):=C(\Psi)$ and $R^{\mathcal{H}}(\theta):=R^{\mathcal{H}}(\Psi)$. This justifies the notation

$$r^{\tau}_{\mathcal{H}}(\theta,\delta):=\sum_{t_i,t_{i+1}\in\tau}\frac{R^{\mathcal{H}}_{t_i}\left(\theta+\delta\mathbf{1}_{(t_i,t_{i+1}]}\right)-R^{\mathcal{H}}_{t_i}(\theta)}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]},$$

where τ is a partition of $[0,T]$.

We now prove the martingale characterization of $(\mathbb{H},\mathbb{F})$-locally risk-minimizing strategies.

Proof of Proposition 3.8. **Step 1.** By using similar arguments to those used in the proof of Lemma 2.2 of [20], first we show that an $(\mathbb{H},\mathbb{F})$-admissible strategy $\Psi=(\theta,\eta)$ with $V_T(\Psi)=\xi$ $\mathbb{P}$-a.s. is $(\mathbb{H},\mathbb{F})$-locally risk-minimizing if and only if Ψ is mean-self-financing and

$$\liminf_{n\to\infty}r^{\tau_n}_{\mathcal{H}}(\theta,\delta)\geq 0,\quad(\mathbb{P}\otimes\langle M\rangle)-\text{a.e. on }\Omega\times[0,T]\tag{A.2}$$

for every bounded $\mathbb{H}$-predictable process δ such that the variation of $\int\delta_u\alpha_u\mathrm{d}\langle M\rangle_u$ is bounded with $\delta_T=0$ and every increasing sequence $(\tau_n)_{n\in\mathbb{N}}$ of partitions of $[0,T]$ tending to identity.
Let $\Psi=(\theta,\eta)$ be an $(\mathbb{H},\mathbb{F})$-admissible mean-self-financing strategy with $V_T(\Psi)=\xi$ $\mathbb{P}$-a.s. such that condition (A.2) is satisfied. Now, take a small perturbation $\Delta=(\delta,\gamma)$ and a partition τ of $[0,T]$. For $t_i,t_{i+1}\in\tau$, we get the following relationship between the $(\mathbb{H},\mathbb{F})$-admissible (but not necessarily mean-self-financing) strategy $\Psi+\Delta|_{(t_i,t_{i+1}]}$ and the $(\mathbb{H},\mathbb{F})$-admissible mean-self-financing strategy associated to $\theta+\delta|_{(t_i,t_{i+1}]}$:

$$r^{\tau}_{\mathcal{H}}(\Psi,\Delta)=r^{\tau}_{\mathcal{H}}(\theta,\delta)+\sum_{t_i,t_{i+1}\in\tau}\frac{\left(\gamma_{t_i}+\mathbb{E}\left[\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\middle|\mathcal{H}_{t_i}\right]\right)^2}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]}.\tag{A.3}$$

Then, by (A.2) it immediately follows that Ψ is $(\mathbb{H},\mathbb{F})$-locally risk-minimizing.
For the converse, let Ψ be an $(\mathbb{H},\mathbb{F})$-locally risk-minimizing strategy. By adapting Lemma 2.1 of [20] to our framework, it is not difficult to show that Ψ is also mean-self-financing. It only remains to prove that condition (A.2) is fulfilled. Let us observe that we may

choose all γ_{t_i} to be 0 in (A.3). By Assumptions 2.8 and 3.2, the following estimates hold:

$$\begin{aligned}\sum_{t_i,t_{i+1}\in\tau} &\frac{\left(\mathbb{E}\left[\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\middle|\mathcal{H}_{t_i}\right]\right)^2}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]}\\ &\leq \bar{K}^2\|\delta\|_\infty\sum_{t_i,t_{i+1}\in\tau}\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\mathcal{H}_{t_i}\right]\mathbf{1}_{(t_i,t_{i+1}]}\\ &\leq \bar{K}^2\|\delta\|_\infty\sum_{t_i,t_{i+1}\in\tau}\rho(t_{i+1}-t_i)\mathbf{1}_{(t_i,t_{i+1}]}.\end{aligned}$$

It is easy to see that this last expression converges to 0 $(\mathbb{P}\otimes\langle M\rangle)$-a.e. on $\Omega\times[0,T]$. Hence, (A.2) is satisfied.

Step 2. We now consider the $\mathbb{F}$-martingale $C(\theta)=(C_t(\theta))_{0\leq t\leq T}$ that represents the cost process associated to an $(\mathbb{H},\mathbb{F})$-admissible mean-self-financing strategy $\Psi=(\theta,\eta)$. Since $C(\theta)$ is square-integrable, we can apply Proposition 2.7 and get the Galthouk-Kunita-Watanabe decomposition of $C_T(\theta)$ with respect to M under partial information, i.e.

$$C_T(\theta)=C_0(\theta)+\int_0^T\mu_u^{\mathcal{H}}\mathrm{d}M_u+O_T,\quad\mathbb{P}-\text{a.s.},\tag{A.4}$$

where $\mu^{\mathcal{H}}\in\mathcal{M}^2_{\mathcal{H}}(0,T)$ and $O\in L^2_{\mathcal{F}}(0,T)$ is weakly orthogonal to M. For a partition τ of $[0,T]$, consider the locally perturbed process associated to the $\mathbb{F}$-martingale $C(\theta)$:

$$C_t\left(\theta+\delta\mathbf{1}_{(t_i,t_{i+1}]}\right)=\mathbb{E}\left[C_T(\theta)-\int_{t_i}^{t_{i+1}}\delta_u\mathrm{d}S_u\middle|\mathcal{F}_t\right],\quad 0\leq t\leq T,\quad 0\leq i\leq N-1.$$

We need the following auxiliary result.

Lemma A.4. *Suppose that Assumptions 2.8 and 3.2 are in force. Then the following statements are equivalent:*

(1) $\liminf_{n\to\infty}r^{\tau_n}_{\mathcal{H}}(\theta,\delta)\geq 0$, $(\mathbb{P}\otimes\langle M\rangle)-$a.e. on $\Omega\times[0,T]$, *for every bounded* $\mathbb{H}$*-predictable process* δ *such that the variation of* $\int\delta_u\alpha_u\mathrm{d}\langle M\rangle_u$ *is bounded with* $\delta_T=0$ *and every increasing sequence* $(\tau_n)_{n\in\mathbb{N}}$ *of partitions of* $[0,T]$ *tending to identity.*
(2) $\mu^{\mathcal{H}}=0$, $(\mathbb{P}\otimes\langle M\rangle)-$a.e. on $\Omega\times[0,T]$, *where* μ^H *is given in* (A.4).
(3) $C(\theta)$ *is weakly orthogonal to* M.

Proof. First we show that the limit in (1) exists $(\mathbb{P}\otimes\langle M\rangle)$-a.e. on $\Omega\times[0,T]$ and equals $\delta^2-2\delta\mu^{\mathcal{H}}$. Similarly to the proof of Proposition 3.1 of [19], consider the difference

$$\begin{aligned}&C_T\left(\theta+\delta\mathbf{1}_{(t_i,t_{i+1}]}\right)-C_{t_i}\left(\theta+\delta\mathbf{1}_{(t_i,t_{i+1}]}\right)\\ &=C_T(\theta)-C_{t_i}(\theta)-\int_{t_i}^{t_{i+1}}\delta_u\mathrm{d}M_u-\left(\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u-\mathbb{E}\left[\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\middle|\mathcal{F}_{t_i}\right]\right).\end{aligned}$$

Then by (A.4) and Lemma 5.4 of [9], we have

$$
\begin{aligned}
&R_{t_i}^{\mathcal{H}}\left(\theta+\delta\mathbf{1}_{(t_i,t_{i+1}]}\right)-R_{t_i}^{\mathcal{H}}(\theta)\\
&=\mathbb{E}\left[\left(C_T\left(\theta+\delta\mathbf{1}_{(t_i,t_{i+1}]}\right)-C_{t_i}\left(\theta+\delta\mathbf{1}_{(t_i,t_{i+1}]}\right)\right)^2\middle|\,\mathcal{H}_{t_i}\right]-\mathbb{E}\left[\left(C_T(\theta)-C_{t_i}(\theta)\right)^2\middle|\,\mathcal{H}_{t_i}\right]\\
&=\mathbb{E}\left[\int_{t_i}^{t_{i+1}}\left(\delta_u^2-2\delta_u\mu_u^{\mathcal{H}}\right)\mathrm{d}\langle M\rangle_u\middle|\,\mathcal{H}_{t_i}\right]+\mathbb{E}\left[\mathrm{Var}\left[\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\middle|\,\mathcal{F}_{t_i}\right]\middle|\,\mathcal{H}_{t_i}\right]\\
&\quad+2\mathbb{E}\left[\mathrm{Cov}\left[\int_{t_i}^{t_{i+1}}\delta_u\mathrm{d}M_u-(C_{t_{i+1}}(\theta)-C_{t_i}(\theta)),\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\middle|\,\mathcal{F}_{t_i}\right]\middle|\,\mathcal{H}_{t_i}\right].
\end{aligned}
$$

Then, this allows to write the quantity $r_{\mathcal{H}}^{\tau_n}(\theta,\delta)$ easily as the sum of three terms. By martingale convergence, the term involving the process $\mu^{\mathcal{H}}$ tends to $\delta^2-2\delta\mu^{\mathcal{H}}$ $(\mathbb{P}\otimes\langle M\rangle)$-a.e. on $\Omega\times[0,T]$, as argued in the proof of Proposition 3.1 of [19]. For the second term, we get the following estimate:

$$
\begin{aligned}
&\sum_{t_i,t_{i+1}\in\tau}\frac{\mathbb{E}\left[\mathrm{Var}\left[\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\middle|\,\mathcal{F}_{t_i}\right]\middle|\,\mathcal{H}_{t_i}\right]}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\,\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]}\\
&\qquad\le\sum_{t_i,t_{i+1}\in\tau}\frac{\mathbb{E}\left[\mathbb{E}\left[\left(\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\right)^2\middle|\,\mathcal{F}_{t_i}\right]\middle|\,\mathcal{H}_{t_i}\right]}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\,\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]}\\
&\qquad=\sum_{t_i,t_{i+1}\in\tau}\frac{\mathbb{E}\left[\left(\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\right)^2\middle|\,\mathcal{H}_{t_i}\right]}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\,\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]}.
\end{aligned}
$$

About the third term, we use the Cauchy-Schwarz inequality for sums and the previous estimate to get

$$
\begin{aligned}
&\left|\sum_{t_i,t_{i+1}\in\tau}\frac{\mathbb{E}\left[\mathrm{Cov}\left[\int_{t_i}^{t_{i+1}}\delta_u\mathrm{d}M_u-(C_{t_{i+1}}(\theta)-C_{t_i}(\theta)),\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\middle|\,\mathcal{F}_{t_i}\right]\middle|\,\mathcal{H}_{t_i}\right]}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\,\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]}\right|\\
&=\left|\sum_{t_i,t_{i+1}\in\tau}\frac{\mathbb{E}\left[\int_{t_i}^{t_{i+1}}\delta_u\mathrm{d}M_u-\left(C_{t_{i+1}}(\theta)-C_{t_i}(\theta)\right)\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\middle|\,\mathcal{H}_{t_i}\right]}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\,\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]}\right|\\
&\le\left(\sum_{t_i,t_{i+1}\in\tau}\frac{\mathbb{E}\left[\left(\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\right)^2\middle|\,\mathcal{H}_{t_i}\right]}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\,\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]}\right)^{\frac{1}{2}}\\
&\quad\cdot\left(\sum_{t_i,t_{i+1}\in\tau}\frac{\mathbb{E}\left[\int_{t_i}^{t_{i+1}}\delta_u^2\mathrm{d}\langle M\rangle_u+\left(\langle C(\theta)\rangle_{t_{i+1}}-\langle C(\theta)\rangle_{t_i}\right)\middle|\,\mathcal{H}_{t_i}\right]}{\mathbb{E}\left[\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\middle|\,\mathcal{H}_{t_i}\right]}\mathbf{1}_{(t_i,t_{i+1}]}\right)^{\frac{1}{2}}.
\end{aligned}
$$

By similar arguments to the ones used in the proof of Proposition 3.1 of [19], it is sufficient to show that

$$\lim_{n\to\infty}\sum_{t_i,t_{i+1}\in\tau}\frac{\left(\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\right)^2}{\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}}\mathbf{1}_{(t_i,t_{i+1}]}=0,\quad(\mathbb{P}\otimes\langle M\rangle)-\text{a.e. on }\Omega\times[0,T],$$

due to Lemma 2.1 of [19]. By Assumptions 2.8 and 3.2, we have

$$\begin{aligned}\sum_{t_i,t_{i+1}\in\tau}\frac{\left(\int_{t_i}^{t_{i+1}}\delta_u\alpha_u\mathrm{d}\langle M\rangle_u\right)^2}{\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}}\mathbf{1}_{(t_i,t_{i+1}]}&\le\bar{K}^2\|\delta\|_\infty^2\sum_{t_i,t_{i+1}\in\tau}\left(\langle M\rangle_{t_{i+1}}-\langle M\rangle_{t_i}\right)\mathbf{1}_{(t_i,t_{i+1}]}\\&\le\bar{K}^2\|\delta\|_\infty\sum_{t_i,t_{i+1}\in\tau}\rho(t_{i+1}-t_i)\mathbf{1}_{(t_i,t_{i+1}]}\end{aligned}$$

and the last expression converges to 0 $(\mathbb{P}\otimes\langle M\rangle)$-a.e. on $\Omega\times[0,T]$.
By (A.4), it is easy to check that $C(\theta)$ is weakly orthogonal to M if and only if $\mu^{\mathcal{H}}=0$ $(\mathbb{P}\otimes\langle M\rangle)$-a.e. on $\Omega\times[0,T]$. It is obvious that (2) implies (1). Since $\lim_{n\to\infty}r_{\mathcal{H}}^{\tau_n}(\theta,\delta)=\delta^2-2\delta\mu^{\mathcal{H}}$ $(\mathbb{P}\otimes\langle M\rangle)-$ a.e. on $\Omega\times[0,T]$, for every bounded $\mathbb{H}$-predictable process δ such that the variation of $\int\delta_u\alpha_u\mathrm{d}\langle M\rangle_u$ is bounded with $\delta_T=0$, to prove that (1) implies (2), for any $\epsilon>0$ and $k>0$ we choose $\delta:=\epsilon\cdot\mathrm{sign}(\mu^{\mathcal{H}})\cdot\mathbf{1}_{\{|\int_0^T\delta_u\alpha_u\mathrm{d}\langle M\rangle_u|\le k\}}$. Clearly, $\int\delta_u\alpha_u\mathrm{d}\langle M\rangle_u$ is bounded and by (1) we deduce that $|\mu^{\mathcal{H}}|\mathbf{1}_{\{|\int_0^T\delta_u\alpha_u\mathrm{d}\langle M\rangle_u|\le k\}}\le\frac{\epsilon}{2}\mathbf{1}_{\{|\int_0^T\delta_u\alpha_u\mathrm{d}\langle M\rangle_u|\le k\}}$, which implies $|\mu^{\mathcal{H}}|\le\frac{\epsilon}{2}$ by letting $k\to\infty$. □

Step 3. Finally, by applying Lemma A.4 we obtain the link between condition (A.2) and the weak orthogonality condition that implies the result. This concludes the proof of Proposition 3.8.

□

Acknowledgements

The third named author was partially supported by the ANR Project MASTERIE 2010 BLAN 0121 01.

References

[1] J.P. Ansel and C. Stricker. Unicité et existence de la loi minimale. In J. Azéma, P.A. Meyer, and M. Yor, editors, *Séminaire de Probabilités XXVII*, volume 1557 of *Lecture Notes in Mathematics*, pages 22–29. Springer, 1993.

[2] F. Biagini and A. Cretarola. Local risk-minimization for defaultable markets. *Mathematical Finance*, 19(4):669–689, 2009.

[3] P. Briand, B. Delyon, and J. Mémin. On the robustness of backward stochastic differential equations. *Stochastic Processes and their Applications*, 97(2):229–253, 2002.

[4] R. Buckdahn. Backward stochastic differential equations driven by a martingale. *Preprint, Humboldt University, Berlin*, 1993.

[5] R. Carbone, B. Ferrario, and M. Santacroce. Backward stochastic differential equations driven by càdlàg martingales. *Theory of Probability and its Applications*, 52(2):304–314, 2008.

[6] C. Ceci. Risk-minimizing hedging for a partially observed high frequency data model. *Stochastics: An International Journal of Probability and Stochastic Processes*, 78(1):13–31, 2006.

[7] C. Ceci. Utility maximization with intermediate consumption under restricted information for jump market models. *International Journal of Theoretical and Applied Finance*, 15(6):24–58, 2012.

[8] C. Ceci. Optimal investment-consumption for partially observed jump-diffusions. In Russo F. Dalang R.C., Dozzi M., editor, *Progress in Probability, Seminar on Stochastic Analysis, Random Fields and Applications VII*. 2013. to appear.

[9] C. Ceci, A. Cretarola, and F. Russo. GKW representation theorem under restricted information. An application to risk-minimization. *Arxiv:1205.3726v2 [math.PR]*, 2012.

[10] N. El Karoui and S. J. Huang. A general result of existence and uniqueness of backward stochastic differential equations. In N. El Karoui and L. Mazliak, editors, *Backward Stochastic Differential Equations*, pages 27–36. Longman, Harlow, 1997.

[11] N. El Karoui, S. Peng, and M. C. Quenez. Backward stochastic differential equations in finance. *Mathematical Finance*, 7(1):1–71, 1997.

[12] H. Föllmer and M. Schweizer. Hedging of contingent claims under incomplete information. In M. H. A. Davis and R. J. Elliott, editors, *Applied Stochastic Analysis*, pages 389–414. Gordon and Breach, 1991.

[13] H. Föllmer and D. Sondermann. Hedging of non-redundant contingent claims. In W. Hildenbrand and A. Mas-Colell, editors, *Contributions to Mathematical Economics*, pages 203–233. 1986.

[14] R. Frey. Risk minimization with incomplete information in a model for high-frequency data. *Mathematical Finance*, 10(2):215–225, 2000.

[15] R. Frey and T. Schmidt. Pricing and hedging of credit derivatives via the innovations approach to nonlinear filtering. *Finance and Stochastics*, 16(1):105–133, 2012.

[16] P. Monat and C. Stricker. Föllmer-schweizer decomposition and mean-variance hedging of general claims. *Annals of Probability*, 23(2):605–628, 1995.

[17] P. Protter. *Stochastic Integration and Differential Equations*, volume 21 of *Applications of Mathematics*. Springer-Verlag Berlin Heidelberg, 2nd edition, 2004.

[18] P. Protter and K. Shimbo. No arbitrage and general semimartingales. *Markov Processes and related Topics: A Festschrift for Thomas G. Kurtz*, 4:267–283, 2008.

[19] M. Schweizer. Risk-minimality and orthogonality of martingales. *Stochastics and Stochastics Reports*, 30(2):123–131, 1990.

[20] M. Schweizer. Option hedging for semimartingales. *Stochastic Processes and their Applications*, 37(2):339–363, 1991.

[21] M. Schweizer. Approximating random variables by stochastic integrals. *Annals of Probability*, 22(3):1536–1575, 1994.

[22] M. Schweizer. Risk-minimizing hedging strategies under restricted information. *Mathematical Finance*, 4(4):327–342, 1994.

[23] M. Schweizer. A guided tour through quadratic hedging approaches. In E. Jouini, J. Cvitanic, and M. Musiela, editors, *Option Pricing, Interest Rates and Risk Management*, pages 538–574. Cambridge University Press, Cambridge, 2001.

[24] P. Tankov. *Pricing and hedging in exponential Lévy models: review of recent results.* Paris-Princeton Lecture Notes in Mathematical Finance. Springer Verlag, 2010.

Claudia Ceci, Dipartimento di Economia, Università degli Studi "G. D'Annunzio" di Chieti-Pescara, Viale Pindaro 42, I-65127 Pescara, Italy.

E-mail address: c.ceci@unich.it

Alessandra Cretarola, Dipartimento di Matematica e Informatica, Università degli Studi di Perugia, via Vanvitelli, 1, I-06123 Perugia, Italy.

E-mail address: alessandra.cretarola@dmi.unipg.it

Francesco Russo, Ecole Nationale Supérieure des Techniques Avancées, ENSTA-ParisTech Unité de Mathématiques appliquées, 828, boulevard des Maréchaux, F-91120 Palaiseau

E-mail address: francesco.russo@ensta-paristech.fr